

# Peer Reviewed

### Title:

Recent Advances in Understanding Flow Dynamics and Transport of Water-Quality Constituents in the Sacramento–San Joaquin River Delta

## Journal Issue:

San Francisco Estuary and Watershed Science, 14(4)

## Author:

<u>Schoellhamer, David H.</u>, U.S. Geological Survey, Portland OR <u>Wright, Scott A.</u>, U.S. Geological Survey, Sacramento, CA <u>Monismith, Stephen G.</u>, Dept. of Civl and Environmental Engineering, Stanford University <u>Bergamaschi, Brian A.</u>, U.S. Geological Survey, Sacramento, CA

#### **Publication Date:**

2016

## Permalink:

http://escholarship.org/uc/item/3vb656d6

#### Acknowledgements:

We thank the Delta Science Program for their support and the Editors of the State of Bay– Delta Science 2016 for their guidance. Richard Norgaard, Michael Healey, Paul Work, and three anonymous reviewers provided constructive comments to previous manuscripts.

#### Keywords:

hydrodynamics, transport, water quality, Sacramento–San Joaquin River Delta, ocean, watershed, anthropogenic

#### Local Identifier(s):

jmie\_sfews\_33444

#### Abstract:

doi: https://doi.org/10.15447/sfews.2016v14iss4art1

This paper, part of the collection of research comprising the State of Bay–Delta Science 2016, describes advances during the past decade in understanding flow dynamics and how waterquality constituents move within California's Sacramento–San Joaquin River Delta (Delta). Waterquality constituents include salinity, heat, oxygen, nutrients, contaminants, organic particles, and inorganic particles. These constituents are affected by water diversions and other human manipulations of flow, and they greatly affect the quantity and quality of benthic, pelagic, and intertidal habitat in the Delta. The Pacific Ocean, the Central Valley watershed, human intervention,



eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.

the atmosphere, and internal biogeochemical processes are all drivers of flow and transport in the Delta. These drivers provide a conceptual framework for presenting recent findings. The tremendous expansion of acoustic and optical instruments deployed in the Delta over the past decade has greatly improved our understanding of how tidal variability affects flow and transport. Sediment is increasingly viewed as a diminishing resource needed to sustain pelagic habitat and tidal marsh, especially as sea level rises. Connections among the watershed, Delta, and San Francisco Bay that have been quantified recently highlight that a landscape view of this system is needed, rather than consideration of each region in isolation. We discuss interactions of multiple drivers and information gaps.

## **Copyright Information:**



Copyright 2016 by the article author(s). This work is made available under the terms of the Creative Commons Attribution <u>4.0 license, http://creativecommons.org/licenses/by/4.0/</u>



eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.



SPECIAL ISSUE: THE STATE OF BAY-DELTA SCIENCE 2016, PART 3

# Recent Advances in Understanding Flow Dynamics and Transport of Water-Quality Constituents in the Sacramento–San Joaquin River Delta

David H. Schoellhamer<sup>\*1</sup>, Scott A. Wright<sup>2</sup>, Stephen G. Monismith<sup>3</sup>, and Brian A. Bergamaschi<sup>4</sup>

#### Volume 14, Issue 4 | Article 1

doi: http://dx.doi.org/10.15447/sfews.2016v14iss4art1

- \* Corresponding author: *dschoell@usgs.gov*
- 1 U.S. Geological Survey Portland, OR 97201 USA
- 2 U.S. Geological Survey Sacramento, CA 95819 USA
- 3 Dept. of Civil and Environmental Engineering Stanford University Stanford, CA 94305 USA
- 4 U.S. Geological Survey Sacramento, CA 95819 USA

# ABSTRACT

This paper, part of the collection of research comprising the State of Bay-Delta Science 2016, describes advances during the past decade in understanding flow dynamics and how water-quality constituents move within California's Sacramento-San Joaquin River Delta (Delta). Water-quality constituents include salinity, heat, oxygen, nutrients, contaminants, organic particles, and inorganic particles. These constituents are affected by water diversions and other human manipulations of flow, and they greatly affect the quantity and quality of benthic, pelagic, and intertidal habitat in the Delta. The Pacific Ocean, the Central Valley watershed, human intervention, the atmosphere, and internal biogeochemical processes are all drivers of flow and transport in the Delta. These drivers provide a conceptual framework for presenting recent findings.

The tremendous expansion of acoustic and optical instruments deployed in the Delta over the past decade has greatly improved our understanding of how tidal variability affects flow and transport. Sediment is increasingly viewed as a diminishing resource needed to sustain pelagic habitat and tidal marsh, especially as sea level rises. Connections among the watershed, Delta, and San Francisco Bay that have been quantified recently highlight that a landscape view of this system is needed, rather than consideration of each region in isolation. We discuss interactions of multiple drivers and information gaps.

# **KEY WORDS**

Hydrodynamics, transport, water quality, Sacramento–San Joaquin River Delta, ocean, watershed, anthropogenic, fluvial forcing

# INTRODUCTION

Movement of water and constituents carried by water within the Sacramento–San Joaquin River Delta (Delta) depend on forcing from the Pacific Ocean, the Central Valley watershed, human intervention, the atmosphere, and internal biogeochemical processes. Semidiurnal and diurnal tides from the ocean slosh water back and forth several kilometers daily through the Delta's complex network of channels. The ocean is a source of salinity, the spatial distribution of which can affect flow dynamics and transport. At a longer time-scale, oceanic sea

level determines sea level rise (SLR) in the Delta. The Central Valley watershed provides freshwater via river flow episodically during California's wet season, and total runoff varies dramatically interannually. The watershed is a source of constituents carried by water including nutrients, contaminants, sediment, and organic particles. Water diversions and associated manipulations of inflow and flow paths alter flow dynamics and transport. Wind can alter Delta water levels and generate wind–waves, which can lift material resting on the bottom into the water column. Biogeochemical processes create sources and sinks of constituents within the Delta. Water and its constituents that exit the Delta in the seaward direction affect San Francisco Bay (the bay).

Technological advances in instrumentation in the past 2 to 3 decades have enabled rapid and continuous measurements of water flow and constituent concentrations. Mike Simpson and Rick Oltmann of the U.S. Geological Survey (USGS) were the first to use acoustic current meters to measure water flow in the Delta, doing so from moving boats (Simpson and Oltmann 1993) and continuously from a fixed station (Oltmann 1995; Ruhl and Simpson 2005). Today, an extensive flow station network in the Delta utilizes acoustic instrumentation (Burau et al. 2016). Optical sensors invented in the 1980s were first deployed in the Delta in the late 1990s to continuously measure suspended-sediment concentration (Wright and Schoellhamer 2005). Optical sensors that continuously measure carbon and nutrients were first deployed in the Delta in the 2000s (Bergamaschi et al. 2012; Downing et al. 2009; Pellerin et al. 2013). When these instruments are deployed with sufficient density to enable several channels or locations to be observed simultaneously, continuous data streams that include all tidal fluctuations can be collected, and these data have led to improved understanding of flow and constituent transport in the Delta.

This paper, part of the State of Bay–Delta Science 2016, summarizes advances in the last 10 years in flow dynamics and how water-quality constituents move within the Delta. Water-quality constituents include salinity, heat, oxygen, nutrients, contaminants, organic particles, and inorganic particles (i.e., sediment). These constituents are affected by water diversions and other human manipulations of flow, and these constituents themselves greatly affect the quantity and quality of benthic, pelagic, and inter-tidal habitat in the Delta. The geographic scope of this paper is the Delta, how the watershed affects the Delta, and how the Delta interacts with the bay and ocean. Though we attempt to cover the breadth of our topic by using a conceptual framework of flow and transport in the Delta, we do not intend to present an exhaustive list of recent publications.

This paper is organized by the primary drivers of flow and transport in the Delta: ocean, fluvial, anthropogenic, atmospheric, and biogeochemical. Our intent is to present new discoveries rather than a conceptual model of flow and transport. Conceptual models available elsewhere include flow (Monsen et al. 2007; Sridharan 2015) and sediment transport (Schoellhamer et al. 2012). Though we do not present a conceptual model, a conceptual framework for the drivers that determine flow and transport within and through the Delta is presented in Figure 1. We also discuss interactions of multiple drivers and information gaps.



**Figure 1** Conceptual framework for drivers of flow and transport in the delta. The black arrows represent flow and transport within and through the delta. Anthropogenic drivers depicted in gray are dams, barriers in the delta, and water exports. Atmospheric drivers depicted in yellow are storms, solar radiation, and wind. Other drivers depicted are biogeochemical processes in green, fluvial inputs in brown, and oceanic exchange in blue.

Other papers that comprise the State of Bay–Delta Science 2016 cover the geographic setting and some specific flow and transport topics in detail, and, thus we minimally include those topics. The geographic setting of the Delta is presented elsewhere in this issue (Healey et al. 2016). Numerical modeling of hydrodynamics and transport in the Delta has greatly advanced in the past 10 years, and is a key tool for testing hypotheses and planning management actions. The numerical modeling paper (MacWilliams et al. 2016) discusses these advances in detail. Movement of fish in the Delta is affected by water flows and is a key management concern. Fish movement is presented elsewhere in this volume (Moyle et al. 2016; Perry et al. 2016).

## FLOW AND TRANSPORT DRIVERS

## **Oceanic Forcing**

The Pacific Ocean, via the bay, affects the Delta by propagation of tides, salinity intrusion, and sea level.

## Tides

The Bay–Delta is a classic example of a coastal plain estuary in which terrestrial freshwater mixes with salt water entering the estuary from the ocean. The primary agent for this mixing is energetic tidal motions (Fischer et al. 1979) forced by propagation of ocean tides into the Bay–Delta (Cheng et al. 1993). At the western end of the Delta, tidal currents are generally 10 to 100 times larger than tidally averaged currents (see Figure 2; Oltmann 1998). Accordingly, in much of the Delta, tides generally cause the instantaneous currents experienced by organisms, sediment erosion, and deposition (Brennan et al. 2002), and vertical turbulent mixing important for example, to phytoplankton dynamics (Jones et al. 2009).

The flow leaving the Delta past Mallard Island and entering the bay is central to regulations of flow and water quality. Direct measurement of Delta outflow has proven to be difficult, so two methods have been used to estimate outflow. The first is the hydrologic balance embodied in Dayflow<sup>1</sup> and the second is to sum the flows measured by four key USGS flow stations (Oltman 1998). Figure 3 compares two flows for water years 2008–2014. Overall, the comparison is quite good, with a slope near 1,  $r^2 = 0.92$ , and an RMS error of 6.4Kcfs. On the other hand, if one examines only low flows, here defined as Dayflow  $\leq 10$ Kcfs, then there is almost no relation between the two flows ( $r^2 = 0.10$ ; RMS error = 5.1Kcfs). Thus, at times when the need is most critical to know what

1. See http://www.water.ca.gov/dayflow/documentation/dayflowDoc. cfm#Introduction







**Figure 3** Delta outflow derived from U.S. Geological Survey (USGS) measurements at four stations (points) as a function of California Department of Water Resources (CDWR) Dayflow estimates of Delta outflow for water years 2008–2014 in thousands of cubic feet per second (Kcfs). The top panel (**A**) is for all flows and the bottom panel (**B**) is for Dayflow values less than 10 Kcfs. Least square linear regression lines are shown in red. For all Dayflow values  $r^2 = 0.92$  and for Dayflow values less than 10 Kcfs  $r^2 = 0.10$ . USGS and CDWR data from *http://cdec.water.ca.gov/*.

outflow is, so as to properly manage project (and gate) operations, the uncertainty is very large.

Tides also provide much of the frictional resistance to the mean tidally averaged flow in the Delta. It is this frictional resistance that ultimately determines patterns of mean flow. For example, even though total freshwater flow toward the pumps is set by pumping rates, the spatial distribution of those flows is controlled mainly by the resistance of the different routes by which freshwater can reach the pumps. Fong et al. (2009) show one way in which this behavior might affect Delta-scale transport: They found that because of bedforms, flows through Threemile Slough experience larger drag for flows from the San Joaquin to the Sacramento than vice versa. This should translate into a stronger tidally averaged flow from the Sacramento to the San Joaquin than would occur were the drag symmetric for flow direction. Consequently, the effect of

asymmetric friction should be to shift a larger fraction of the overall outflow through Threemile Slough rather than through more northern channels such as Georgiana Slough.

Parameters that quantify a physical, chemical, or biological constituent of the aquatic environment may be misrepresented by low frequency data that fail to observe tidal variations (Lucas et al. 2006). To better understand biogeochemical processes, newly developed optical instruments of various types have been deployed in the Delta and upstream. The new instrumentation can make unattended measurements *in situ* fairly frequently (e.g., every 15 minutes). This capability is particularly important in a tidal setting where tidal advection and dispersion continuously affect constituent concentrations. For example, Ganju et al. (2005) and Downing et al. (2009) used data from Browns Island to assess the minimum sampling frequency necessary to accurately calculate advective and dispersive fluxes in a tidal wetland. That frequency was on the order of three per hour.

The shape of the tidal wave changes as it propagates upstream into the Delta, which creates tidal asymmetry and retains sediment in some parts of the Delta. Morgan-King and Schoellhamer (2013) found an estuarine turbidity maximum in the backwater Cache Slough complex created by tidal asymmetry (peak flood tide currents are greater than peak ebb tide currents), a limited tidal excursion, and windwave re-suspension. During the study, there was a net export of sediment, though sediment accumulates within the region from landward tidal transport during the dry season. Sediment is continually re-suspended by both wind-waves and flood tide currents. The suspended-sediment mass oscillates within the region until winter freshwater flow pulses flush it seaward. The hydrodynamic characteristics within the backwater region–such as low freshwater flow during the dry season, flood tide dominance, and a limited tidal excursion-favor sediment retention. This sediment retention is reflected in the relatively fine bed sediment found in the Cache Slough area in comparison to other regions of the Delta (Marineau and Wright, forthcoming). Relatively high turbidity makes the Cache Slough complex favorable habitat for Delta Smelt (Nobriga et al. 2005). These isolated backwater regions used to be common in the Delta (Whipple et al. 2012) but are now rare because channels in much of the Delta are now interconnected to convey water to pumps or to convey floodwaters to the ocean. Restoration of isolated dead-end channels may help ecosystem restoration.

A concept that has evolved during the last decade is that transport in the Delta is driven more by tidal flow than tidally averaged flow. A longstanding conceptual model of transport in the Delta neglects tidal motions and dispersion; i.e., transport of heat, pollutants and non-motile organisms is determined entirely by advection by subtidal flows. For example, according to this model, mean flows along the Old and Middle River corridor act (among other things) to transport small Delta Smelt to the export pumps, such that restrictions on Old and Middle River flows can be used to limit entrainment (Kimmerer 2008). The basis for this assumption is that dispersion in the tidal channels is likely to be relatively weak

(Fischer et al. 1979; Ho et al. 2008), and that the effects of tidal motions average out, leaving net motions as the only mechanism for net advection. On the other hand, studies of dispersion in the Delta using numerical models show that dispersion associated with tidal flows through the many junctions of the Delta can be substantial (1995 personal communication between E. List and SGM, unreferenced, see "Notes"; Monsen 2000; Sridharan 2015). The reason for this is that splitting flows at junctions drives what Ridderinkhof and Zimmerman (1992) describe as "chaotic mixing." Observational evidence for this behavior was provided by Monismith et al. (2009) who suggested that the large heat fluxes out of the San Joaquin system required to close the overall heat balance could be accounted for with remarkably large dispersion coefficients such as those that typify chaotic mixing.

The practical consequence of strong Delta-scale dispersion is that overall transport of constituents through the system may differ significantly from what might be expected in the absence of dispersion; in particular, patterns and rates of organism entrainment at the export pumps might not be simply related to tidally averaged flow patterns. Thus, numerical models of transport used to inform policy must correctly represent the mixing effects of junctions. The challenge of this task is demonstrated by the complex flow behavior seen at the Mokelumne River-Georgiana Slough junction described by Gleichauf et al. (2014) (see also Gleichauf 2015). Tidal flows through this junction produce fronts (regions of separation) and secondary flows, both of which can significantly affect the paths taken by materials and organisms that pass through the junction (Figure 4). Nonetheless, as seen in modeling presented in Wolfram et al. (2016), on average, tidally varying flow trajectories produced by a well-resolved, 2-D, depth-averaged circulation model agreed well with the results of a high-resolution, non-hydrostatic 3-D model, suggesting that practical 2-D models might provide sufficiently accurate representations of junction flows to calculate large-scale transport. In contrast, 1-D models in common use (e.g., Kimmerer and Nobriga 2008) that assume complete mixing at junctions can significantly err in predictions of large-scale dispersion in the Delta, as well as in the



**Figure 4** Drifter paths (A) and SUNTANS streamlines (B) reveal a recirculation zone at the west junction of Georgiana Slough (GS) and the Mokelumne River (MOK). Source: Gleichauf et al. (2014).

timing and rates of entrainment at the export pumps (Sridharan 2015).

## Salinity

Freshwater flow affects the position of the lowsalinity zone (LSZ) in northern San Francisco Bay, which is important habitat for many fish species (Kimmerer et al. 2013). The landward extent of salt from the ocean is determined by freshwater flow (Q) into the estuary. To control the landward extent of salinity, regulations require that Q be adjusted to achieve a desired distance a salinity of 2 is from the Golden Gate Bridge (X2) (Jassby et al. 1995). Given how strongly X2 regulations affect water supply, forecasting how X2 depends on Q is an important

component of project operations. Analysis of the flow and salinity data from 1967 to 1991 (Jassby et al. 1995; Monismith et al. 2002) suggested that the time-scale of X2 response to flow is approximately 2 weeks, and that at steady flow X2 is proportional to  $0^{-1/7}$ -behavior that Monismith et al. (2002) suggest reflects the effects of tidally varying stratification on turbulent mixing, and hence the transport of salt by gravitational circulation. Using a synthesis of numerical modeling and analysis of more recent observations, MacWilliams et al. (2015) argue that: (1) the response of X2 is fast when Q is large, and slow when Q is small; and (2) the dependence of X2 on Q is  $Q^{-1/4}$  – behavior that would be predicted from the classical theory of gravitational circulation accounting for actual variations in width and depth of the estuary. MacWilliams et al. (2015) also point out the central challenge of predicting X2 when Q is small: during lowflow conditions, uncertainty in Q can be larger than Q itself. They advocate using a numerical model to define the low-flow X2-0 relation and then (if necessary) using observed X2 variations to infer flow.

Note that both the Jassby et al. (1995) and MacWilliams et al. (2015) models imply symmetrical responses of X2 to flow increases and decreases. In contrast, Chen (2015) showed that the response to flow increases is much faster than the response to flow decreases, in essence reflecting that response time depends on the state of the estuary (X2) as well as on the flow. Monismith (in press) used this idea to construct and validate an X2 model in which the rate of change of X2 is proportional to the difference between the current value of X2 and the equilibrium value of X2 based on the current value of Q. Since this latter model is based on the governing equations, rather than being purely empirical, it may be viewed as preferable from a theoretical standpoint. On the other hand, in practical terms, the physics-based model has guite similar accuracy to the timeseries model (Figure 5). That said, one practical consequence of the asymmetrical model is that it implies that pulsed flows require a larger volume of water to maintain a given average value of X2 than does the steady flow that corresponds to that average value of X2.

Observations of salinity in the main channel of northern San Francisco Bay show that depth-

#### DECEMBER 2016



**Figure 5** X2 for 6 water years derived from USGS–USBR–CDWR fixed station CTD data (solid grey line) and as predicted by the MacWilliams et al. (2015) time series model (dashed line) and the Monismith (forthcoming) model (solid black line)

averaged salinity depends primarily on the distance of a given point in the estuary relative to X2 (Monismith et al. 2002). Measurements of X2 during the fall of 2011 (Stacey et al., in prep.) show that salinities in Honker and Grizzly bays tend to be lower in general than in the main channel, and respond more slowly to changes in flow and spring– neap variations in tidal mixing that also produce variations in X2 (Monismith, forthcoming).

## Sea Level Rise

In addition to tides and salt, future sea level rise (SLR) will propagate into the Delta. Both the salinity field and tidal motions are the result of interaction of the Bay–Delta with physical forcing that itself varies in time. Outflow of freshwater from the Delta, the key

factor that suppresses upstream propagation of salt, will likely change in the future because of changes in precipitation patterns caused by anthropogenic climate change (Cloern et al. 2011). At the same time, SLR is expected to have two effects: (1) gravitational circulation will be stronger when channel depths are larger, increasing upstream dispersive salt fluxes so the outflow required to maintain a particular value of X2 will be larger than it is now (Chua and Xu 2014); and (2) because of the likely flooding of large flat areas adjacent to the current Bav-Delta. frictional damping of tides will be stronger, and so tidal propagation through the system will likely change (Holleman and Stacey 2014). We note that this latter effect has not yet been investigated for the Delta itself-Holleman and Stacey's modeling did not include the Delta. Sea level rise will also influence

water diversions. Wang et al. (2011) simulated future water diversions from the Delta and found that at the end of the 21st century salinity intrusion from SLR of 61 cm will reduce the amount of freshwater available for water diversions.

Swanson et al. (2015) developed a Delta marsh elevation model and found that the magnitude of SLR over the next century was the primary driver of marsh surface elevation change. They defined marsh accretion parameters to encapsulate the range of observed values over historic and modern timescales, based on measurements from four marshes in high- and low-energy fluvial environments. In addition, they modeled possible future trends in sediment supply and mean sea level. They also conducted a sensitivity analysis of 450 simulations that encompassed a range of porosity values, initial elevations, organic and inorganic matter accumulation rates, and SLR rates. More than 84% of the scenarios resulted in sustainable marshes with a moderate 88 cm of SLR by 2100, but only 32% and 11% of the scenarios resulted in surviving marshes when SLR was increased to 133 cm and an upper bound of 179 cm, respectively. Sediment supply was the next most important controlling factor.

## **FLUVIAL FORCING**

The Sacramento River is the primary source of freshwater flow and sediment to the Delta. Wright and Schoellhamer (2005) found that, for the period from 1999 to 2002, the Sacramento River supplied approximately 85% of the water and sediment to the Delta. The second-largest source, the San Joaquin River, supplied about 12%, with the remaining supply from smaller tributaries such as the Cosumnes and Mokelumne rivers. The Sacramento and San Joaquin rivers both have USGS stream gages that have been in operation since 1957 measuring daily discharge and suspended sediment load (114447650 Sacramento River at Freeport and 11303500 San Joaquin River near Vernalis). Over this timeperiod, the average discharge and sediment loads on the Sacramento River were about 23,000 ft<sup>3</sup>/s and 5,400 tons/day (note: this does not include the Yolo Bypass); on the San Joaquin River, these averages were about 4,800 ft<sup>3</sup>/s and 900 tons/day. This suggests that the period from 1999 to 2002

roughly represents at the long-term averages, and that the Sacramento River supplies about 85% of the freshwater flow and sediment to the Delta. The majority of the sediment supply occurs during the winter wet season, typically between December and April. During 1999 to 2002. Wright and Schoellhamer (2005) calculated that 82% of the sediment supply occurred during the wet season, which constituted 31% of the total time Climate change will alter future discharge to the estuary. Cloern et al. (2011) simulated two warming scenarios, one with curtailed greenhouse gas emissions and one with continually increasing emissions. The scenario with curtailed emissions featured statistically insignificant decreases in precipitation and runoff from 2010 to 2099; the increasing emission scenario had statistically significant decreases in both. Both scenarios had statistically significant decreases in percentage of runoff that was snowmelt.

## Dams

Dams and reservoirs in the watersheds draining to the Delta have affected the timing and magnitude of freshwater flows and sediment loads. In particular, sediment trapping in reservoirs has strongly affected sediment supply to the Delta. Reservoirs in the Delta watershed capture and store rainfall and snowmelt runoff for a variety of purposes, leading to changes in the flow hydrograph downstream from the dam. These changes typically take the form of decreased peak flows and elevated flows during other times of the year (Singer 2007). However, extreme flooding events still occur in the watershed (for example, 1986, 1997) because of areas of unregulated runoff as well as spills from reservoirs during periods of very high inflow. These extreme events are flood-managed in the valley, downstream from the reservoirs, by a series of weirs and bypass channels (such as Yolo and Sutter bypasses, Singer 2007; Singer and Aalto 2008).

The dams affect sediment supply more than flow, because the larger reservoirs capture approximately 100% of the incoming sediment load, releasing clear water and thus decreasing the downstream sediment supply. The two largest reservoirs in the Sacramento River watershed, Shasta Lake on the upper Sacramento River and Lake Oroville on the Feather River, capture runoff and sediment from large

mountainous areas that were likely large sources of sediment to the Delta pre-dam (some portion was also likely deposited in the Valley). Reservoir sedimentation is one factor that contributes to the approximately 50% decline in Sacramento River sediment loads since stream gaging began in 1957 (Wright and Schoellhamer 2004). In the present condition, the primary sources of sediment to the Valley floor and Delta are the largely unregulated Sacramento River tributaries in the northern Valley, that enter the Sacramento River downstream from Shasta Dam (such as Cottonwood Creek, Cow Creek, and Elder Creek). Singer and Dunne (2001), as well as analyses of unpublished data from recent turbidity monitoring along the mainstem Sacramento, indicate that sediment runoff is generated in these areas during intense rainfall events, and transported down the Valley by the Sacramento River to the Delta (with some deposition along the way, Singer and Dunne 2001; Singer and Aalto 2008). The Feather-Yuba-Bear watersheds contribute sediment as well: however, it is likely reduced from pre-dam sediment supply. The current-day supply of sediment from the Feather-Yuba-Bear watersheds is a data gap that hinders definitive analysis of sediment sources. That said, the primary source of sediment to the Delta today is likely the Sacramento River watershed between Shasta Dam and the Butte City area.

# First Flush

The first major runoff event of the wet season, which typically occurs in the late fall or early winter, is commonly referred to as the "first flush," and has particular significance in the Delta. For sediment transport, this event typically has the highest suspended-sediment concentrations, and results in a rapid change in turbidity throughout the Delta (Figure 6, Wright and Morgan 2015). The underlying mechanism responsible for this is sediment supply limitation. During dry periods, landscape disturbances increase the supply of sediment available for transport during runoff events. As the wet season progresses, sediment supply is gradually depleted, leading to decreased peak sediment concentrations (for the same discharge).

Notably, first flush sediments are typically high in associated contaminants and organic carbon (Bergamaschi et al. 2001; Hladik et al. 2009). Hladik et al. (2009) noted that suspended contaminant concentration was highest proximate to agricultural areas, but tended to be lower in the mainstem of the San Joaquin River. Likewise, toxicity was found to be higher in agricultural areas (Weston et al. 2008). Weston et al. (2014) studied pesticides in Cache Slough and found no pyrethroids during the dry season, but pyrethroids were toxic to a test organism after storms. Urban and some agricultural runoff into Ulatis Creek, a tributary of Cache Slough, was responsible.

There is also concern about the ability of sediments to transport mercury to the Delta during first flush. Elevated concentrations of mercury in suspended sediments have been reported (Domagalski et al. 2004; Roth et al. 2001). The concern is that these sediments will be deposited in reducing environments such as wetlands (Bergamaschi et al. 2012) where the mercury will be transformed into its organic and more toxic form: methylmercury (Bergamaschi et al. 2011).

Increases in Delta turbidity during the first flush event have been linked to Delta Smelt habitat, migration, and entrainment at export pumps (Moyle et al. 2016). During first flush in the lower Sacramento River, Delta Smelt migrate landward using lateral differences in tidal currents and turbidity (Bennett and Burau 2015).

During storms, when order-of-magnitude changes in concentration can occur in a matter of minutes, high-frequency sampling is necessary. Importantly, these variations are not necessarily in concert with variation in other parameters such as total suspended sediment. Saraceno et al. (2009), working in Willow Slough, a tributary of the Sacramento River, found that the peak in dissolved organic carbon associated with the storm did not align with either the peak in the hydrograph or the peak in TSS, consistent with their different sources (Florsheim et al. 2011; Oh et al. 2013).

Even for steady flows in rivers, sampling at high frequency may be necessary for chemical constituents with multiple sources or ones that are reactive in the environment. Pellerin et al. (2009) showed that nitrate loads calculated in the San Joaquin River using traditional methods can under- or overestimate the true load by more than 30% because of



**Figure 6** First flush along the mainstem of the Sacramento River in 2012. Turbidity (A), tidally-averaged discharge (B), and velocity (C). Turbidity varies with discharge at Freeport upstream from the Delta. Dispersion decreases the peak turbidity in the downstream direction. Travel time and bidirectional flow in the Delta delay the arrival of turbidity at Rio Vista and Mallard Island. Source: Wright and Morgan (2015).

variation in the nitrate concentration. A sampling frequency of approximately four per hour was necessary to accurately calculate loads and resolve sources.

# Sedimentation

By definition, the Delta is a depositional environment for sediment. Deposition of sediment and organic matter contributes to building tidal wetlands in the Delta since the most recent sea level lowstand. The modern sediment budget of the Delta was evaluated by Wright and Schoellhamer (2005) and Morgan-King and Wright (2015), based on high-frequency monitoring of suspended sediment loads at a network of stations in the Delta. For the period from 1999 to 2002, Wright and Schoellhamer (2005) computed the total incoming suspended sediment load to the Delta to be 6.6  $\pm$ 0.9 million metric tons (Mt) and the total outgoing sediment load to be 2.2  $\pm$ 0.7 Mt, resulting in 4.4 ±1.1 Mt of sediment deposition over the 4-year period (about 67% of the incoming load). Bed load is approximately two orders of magnitude less than suspended load (Schoellhamer et al. 2012). The magnitude of deposition in the Delta can also be roughly evaluated over a longer time-period. McKee et al. (2013) computed sediment loads at Mallard Island (the downstream boundary of the Delta) for the period from 1995 to 2010, and incoming sediment loads are also available for this time-period for the Sacramento River (USGS Freeport gage) and San Joaquin River (USGS Vernalis gage). The missing piece of information for this time-period is the sediment load coming from the Yolo Bypass, which was found by Wright and Schoellhamer (2005) by using data from 1957 to 1961 and 1980 to be about 28% of the load at Freeport for 1999 to 2002. If the Yolo Bypass inputs are ignored, the average magnitude of deposition from 1995 to 2010 was 60%; if Yolo Bypass inputs are assumed to be 28% of Freeport loads, this magnitude increases to 67%. This suggests that the modern, decadal-scale sediment budget of the Delta is such that about 60% to 70% of the incoming suspended sediment is deposited within the Delta. This percentage varied from year to year, with a low of 39% in water year 1995 (a wet year, 1.63 Mt of deposition) and a high of 88% in Water Year 2007 (a dry year, 0.91 Mt of deposition). From 1995 to 2010, Water Year 1996 had the greatest

depositional mass (4.53 Mt, 82%) and Water Year 2008 the least (0.27 Mt, 56%).

Morgan-King and Wright (2015) used a more extensive monitoring program to evaluate sediment budgets within the Delta for 2011 to 2013 for specific regions: the north Delta, central Delta, and south Delta. In general, the south Delta tends to have the highest trap efficiency (fraction of incoming sediment that is deposited), because the south Delta tends to be dominated by tidal forcing. In contrast, the north Delta tends to have the lowest trap efficiency, even though the incoming sediment load to the north Delta is by far the highest. This is because the Sacramento River can be dominated by fluvial forcing during high flows such that large amounts of sediment are passed through this region under these conditions. The central Delta tends to experience the most variability in deposition, primarily depending on upstream conditions in the Sacramento and San Joaquin rivers. During high flows, sediment loads into the central Delta through Georgiana Slough and from the San Joaquin River can be large, resulting in high deposition rates. In 2011, which was a relatively wet year, about 50% of the sediment that entered the central Delta was deposited (the highest among the three regions), whereas in 2012, which was a relatively dry year, only about 23% of the incoming sediment was deposited. This illustrates that the central Delta sediment budget depends on the upstream sediment supply.

Particles are not limited to inorganic sediment, and include plankton. Turbulence affects whether particles will remain suspended and susceptible to transport. Thus, turbulence affects the time plankton remain productive in the photic zone and whether this material will be intercepted by benthic grazers such as clams (Lehman et al. 2009; Lucas and Thompson 2012).

Freshwater flow from the Delta enters and affects San Francisco Bay. Larger Delta outflow can freshen central San Francisco Bay, and the resulting density difference flushes south San Francisco Bay, exporting sediment (Figure 7, Shellenbarger et al. 2013). When Delta outflow is small, central San Francisco Bay remains salty, there is little density difference between the central and south bays, and tidally averaged sediment transport is landward. Sediment



**Figure 7** Springtime monthly seaward sediment flux at the Dumbarton Bridge as a function of Delta outflow. Data are for April, May, and June 2009–2011 (Shellenbarger et al. 2013)

supply for the South Bay Salt Pond Restoration Project, the largest tidal wetland restoration project on the west coast of the United States, depends on salinity conditions created by Delta outflow. This demonstrates the dependence of south bay flushing and water quality on Delta outflow, first recognized by McCulloch et al. (1970).

Improved estimates of the historical supply of freshwater and sediment from the Central Valley watershed to the Delta and bay have been developed. Moftakhari et al. (2013) used tidal water level data recorded at San Francisco and tidal theory on the interaction of tides and river discharge to estimate discharge from 1858 to 1929. They found that the annual flow is now 30% less than before 1900, and confirmed that the flood of January 1862 was the largest since 1858. Moftakhari et al. (2015) used measurements of Sacramento River water level to extend this discharge record back to 1849. They also used historical sedimentation data to estimate a timeseries of sediment supply for 1849 to 1929. About 55% of the sediment delivered to the estuary between 1849 and 2011 was the result of anthropogenic alteration in the watershed that increased sediment supply. Hydraulic mining in the Central Valley watershed created an initial pulse of sediment and,

subsequently, urbanization in the San Francisco Bay Area increased agricultural land use, and forestry practices may have increased sediment supply. Sediment supply decreased about 50% since the 19th century, and the fraction of sediment delivered during winter has increased while the fraction delivered during spring has decreased, mimicking discharge patterns.

Since about 1900, the Central Valley watershed and Delta appear to be geomorphically adjusting as sediment supply from the watershed has decreased (Schoellhamer et al. 2013). Hydraulic mining from the mid-1850s to the mid-1880s created a pulse of sediment and aggradation, which was followed by a period of decreasing sediment supply and degradation. The period of adjustment from decreasing sediment supply may have ended about 1999, resulting in a stable regime. Hestir et al. (2013) analyzed monthly total suspended solids concentration data from the delta from 1975 to 2010 and found that large floods in 1983 and 1998 caused step decreases, likely from removal of erodible sediment (Figure 8). From 1999 to 2010 there was a significant decreasing trend in total suspended solids in the Delta, possibly resulting from supply limitation or trapping by aquatic vegetation. Hestir

#### DECEMBER 2016



**Figure 8** Mean total suspended solids concentration (TSS) in milligrams per liter (mg L<sup>-1</sup>) from 6 monthly sampling stations in the Delta, 1975–2010, from Hestir et al. (2013). Dates corresponding to a significant break point are indicated with a vertical dashed line. The 95% confidence interval around the break point is represented by the shaded gray area. The solid lines indicate the trends for the periods separated by breakpoints. Before the first breakpoint and after the second decreasing trends were statistically significant (p < 0.05). For the period between breakpoints the trend is graphically represented by the mean. Data from *http://www.water.ca.gov/bdma/meta/discrete.cfm* 

et al. (2016) found that Delta sites with the greatest cover of submerged aquatic vegetation (SAV) had the greatest decreasing trend of total suspended solids. Schoellhamer et al. (2013) hypothesize that it is likely that the estuary and watershed can still adjust, but further adjustment will be as steps that occur only during greater floods than previously experienced during the adjustment period.

If more precipitation falls as rain rather than snow, creating greater runoff in winter and less in spring, as is the trend (Moftakhari et al. 2015); if storm intensity continues to increase (Russo et al. 2013); if reservoir trapping efficiency does not change; and if there is no more post-hydraulic mining geomorphic adjustment (i.e., erodibility of the watershed does not change, Schoellhamer et al. 2013), then sediment supply to the Delta may increase during the 21st century.

## Water Temperature

Water temperatures in the Delta are generally determined by surface heat fluxes (Monismith et al. 2009; Gleichauf 2015) and thus can be well correlated with atmospheric temperatures (Wagner et al. 2011). Inflow temperatures are generally lower than temperatures in the interior of the Delta (Figure 9), so inflows can affect the spatial gradient of water temperature. Wagner et al. argued that any effect of inflows on the overall temperature was only discernible on short time-scales (approximately 1 month or less). On the other hand, large horizontal tidally averaged heat fluxes (of the same scale as diurnally averaged heating in summer) attributed to dispersion were found to be necessary to close the heat balances in the Sacramento Deep Water Ship Channel (SDWSC) at the Port of Stockton (Monismith et al. 2009) and the Delta as a whole (Gleichauf 2015), suggesting that flow-based heat transport may in fact be an important determinant of Delta temperatures.

An interesting aspect of the temperature distribution in 2011 was the seeming importance of cold water entering the Delta from the Merced, Tuolumne, and Stanislaus rivers, i.e., inflows much closer to the Delta than the more southerly sources of the San Joaquin (e.g., Friant Dam). From a management perspective, this suggests that cold water pools in reservoirs closer to the Delta might be useful to



**Figure 9** Delta wide average summer temperatures for 2011 and 2014. Source: Gleichauf (2015).

help manage temperatures in the Delta, although this would also require managing the temperature structure in those reservoirs through the judicious use of selective withdrawal (Anohin et al. 2006; Fischer et al. 1979).

# **ANTHROPOGENIC FORCING**

Anthropogenic forcing of the Delta takes a variety of forms, such as (1) changes in volume and timing of Delta inflows (Moftakhari et al. 2013); (2) reductions in sediment fluxes into the Delta (Wright and Schoellhamer 2004); and (3) modification of mean and tidal flows through the operation of pumps, gates, and barriers. We presented the first two topics earlier in this paper; we discuss the later topic below.

Near the upstream end of tidal influence (around Freeport), the Sacramento River divides into several distributary channels, to the east and west of the mainstem. The east-side distributary channel, Georgiana Slough, connects with the lower Mokelumne River and, ultimately, the San Joaquin River in the central Delta. Along with the Delta Cross Channel (DCC), which connects the Sacramento River to the Mokelumne system when its gates are open, Georgiana Slough is the primary pathway for water and sediment to move from the Sacramento River to the central Delta. As shown by Wright and Morgan (2015), this pathway can result in elevated turbidity and sediment concentration in the south Delta, because of the landward net flows in Old and Middle rivers driven by pumping facilities in the south Delta. Once sediment enters the central Delta through the Georgiana Slough pathway, it is gradually advected southward though Old and Middle rivers into the south Delta; Wright and Morgan (2015) calculated a travel time of about 2 weeks from Freeport to the south Delta for a first flush sediment pulse in December 2012. This slow advection of sediment southward, which increases turbidity in the south Delta, has important implications because elevated turbidity in the region of the pumping facilities has been linked to high entrainment of Delta Smelt at the facilities (Grimaldo et al. 2009), which can lead to curtailments of federal and state water deliveries.

Both observations (Gleichauf et al. 2014) and numerical modeling of the Delta (Monsen et al.

2007) show that gate operations significantly affect hydrodynamics and transport patterns in the Delta. When the DCC gate is closed, Sacramento River water reaches the pumps through Threemile Slough and through the confluence region (Monsen 2000; Monsen et al. 2007). Because these transport paths are closer to the higher-salinity waters of the bay, salinities in the western Delta tend to be higher when the DCC is closed than when it is open, so meeting water quality standards in the Delta may require greater outflows into the bay. In a similar fashion, Monsen et al. (2007) show that operation of the Head of Old River Barrier (HORB) may significantly affect residence time in the SDWSC: residence time for particles in the SDWSC were five times smaller when the HORB was in place than when it was absent, a difference that may be important to the formation of hypoxic waters in that reach of the San Joaquin. Finally, it seems likely that tidally operated gates, notably the one in Montezuma Slough, can rectify tidal forcing and thus have significant effects on transport and salinity as well. For example, C. Enright (unpublished manuscript) suggests that when the Montezuma Slough gate is being operated to tidally pump freshwater into Suisun Marsh, it effectively diverts approximately 2500 cfs, possibly a large fraction of the dry-weather Delta outflow (approximately 3,000 to 8,000 cfs) that might otherwise be passing through Suisun Bay.

# **ATMOSPHERIC FORCING**

The water surface is the interface between the atmosphere and Delta waters. Wind can alter Delta water levels and generate wind-generated waves, which can lift material resting on the bottom into the water column. The Delta breeze is the dry season afternoon and evening wind that blows from cooler Suisun Bay through the Delta and into the warmer Central Valley. Heat exchange at the water surface warms or cools Delta waters.

Wind-generated waves enhance sediment re-suspension in some shallow parts of the Delta. Ganju et al. (2005) found that wind wave re-suspension adjacent to Browns Island contributed suspended sediment to the main tidal channel on the island. Morgan–King and Schoellhamer (2013) observed increased suspended-sediment concentrations (SSC) in Cache Slough and the SDWSC (which has a substantial fraction of shallow water) during spring and summer. These increases in SSC were well-correlated with wind speed. Landward sediment flux was greatest during these windy seasons, indicating that sediment retention was enhanced.

Along with wind wave re-suspension of sediments, wind waves also can greatly enhance mixing from wave breaking (Jones and Monismith 2008), an effect in the Delta that is likely to be especially pronounced in the shallow, open-water regions of Franks Tract (Jones et al. 2008) and, presumably, Mildred Island. The enhancement of near-surface turbulence by wave breaking is important in that strong turbulent shears and energetic vertical mixing, both effects of wave breaking, can act to break up colonies of cyanobacteria such as Microcystis (O'Brien et al. 2004) and, ultimately, suppress blooms (c.f. Huisman et al. 1999). More generally, wind mixing is important to phytoplankton dynamics in that nearsurface turbulence plays a big role in determining the light climate experienced by individual phytoplankton cells (MacIntyre 1993).

Intensity of rainfall increased from 1890 to 2010 (Russo et al. 2013). Data from over 600 precipitation stations in the greater San Francisco Bay area, including the Delta and part of the Central Valley, were analyzed. The intensity of the largest (less frequent) storms increased the most, and a greater fraction of rain fell during large events. More intense storms increase the steepness of runoff hydrographs and increase sediment transport. They may also increase the release of metals from inter-tidal sediments (Moskalski et al. 2013).

Climate change will warm Delta waters. Delta water temperatures are strongly related to air temperature, and not as much to Delta inflow rates (Monismith et al. 2009), so increased air temperatures will warm Delta waters. For scenarios of curtailed and increasing greenhouse gas emissions, Cloern et al. (2011) and Wagner et al. (2011) estimate there will be statistically significant increases in Sacramento River and Delta water temperatures. Delta water temperatures will increases up to 5° C in summer, a change that may prove lethal for Delta Smelt. Given that growth rates for cyanobacteria increase exponentially with temperature (Robarts and Zohary 1987), this temperature rise may also enhance the likelihood of cyanobacterial blooms (Paerl and Huisman 2009). Whether this warming may increase thermal stratification was not evaluated.

Measurements made in the summers of 2004 and 2005 in the SDWSC of vertically varying temperatures show the formation and destruction of diurnal stratification that can be as strong as 2°C from the top to the bottom of the water column (Figure 10, Monismith et al. 2008). This temperature stratification does appear to be dynamically significant, and is the result of turbidity-dependent solar heating and mixing provided by the wind and surface cooling, as well as by turbulence production near the bottom. The practical significance of this stratification is that, given that existing descriptions of the temperature field of the Delta are based on the existing network of sensors, which are largely nearsurface, it may be necessary to reconsider the nature of the effects of long-term temperature variability on fish habitat (Wagner et al. 2011; Gleichauf 2015). In particular, it may be possible that there are daytime refugia of colder water that temperature-sensitive fish such as Delta Smelt can use to avoid warmer surface waters. It would be most useful if the existing temperature sensor network were expanded to include near-bottom as well as near-surface temperatures.

# **BIOGEOCHEMICAL FORCING**

Biogeochemical processes within the Delta can alter dissolved substances and suspended particulates. For example, Alpers et al. (2014) studied methyl mercury production in the Yolo Bypass and found methylation produced the highest concentrations in drainage from wild rice fields during harvest, and in white rice fields with decomposing rice straw during flooding. An example of biota affecting transport is invasive SAV in Delta channels. SAV began increasing significantly in the 1970s, and sites that now have greater SAV coverage had a greater turbidity decline in the later 20th century (Hestir et al. 2016). SAV can slow water movement and reduce bed shear stress, promoting sedimentation and reducing SSC. Turbidity trends were corrected for the declining sediment



**Figure 10** Diurnal variation in temperature stratification at four stations in the Stockton Deep Water Ship Channel. The stations are ordered in the downstream direction and specific locations are given in Monismith et al. (2009). At each station, temperatures were measured at ca. 2m intervals in the vertical.

supply (Wright and Schoellhamer 2004) using SSC data immediately upstream from the Delta. Hestir et al. (2016) estimate that 21% to 70% of the total declining turbidity trend results from SAV expansion.

## DISCUSSION

The *State of Bay–Delta Science 2008* (Healey et al. 2008) states:

... it is now recognized that the twice-daily tides are also extremely influential, causing powerful flow reversals through much of the Delta that amplifies dispersive mixing in both directions. Geometric features of Delta waterways, such as bends, junctions, shallow water areas and levees, all influence water transport and residence times.

Appreciation and consideration of tidal transport and effects were once limited to physical studies, but it has become more common in water quality and ecological studies, and has helped to improve our understanding of these fields (e.g., Bergamaschi et al. 2011; Bennett and Burau 2015).

## **Interactions of Multiple Drivers**

We have summarized the state of the science for flow and transport by the primary drivers: ocean, fluvial, anthropogenic, atmospheric, and biogeochemical (Figure 1). Flow and transport in the Delta are determined by the interactions of these drivers, and management decisions must consider all these drivers.

Drought reduces the effect of fluvial forcing in the Delta and increases the effect of ocean forcing. The drought that began in 2012 decreases sediment supply from the watershed, increases residence times in the Delta, can increase salinity depending on regulation by water managers, and reduces hydrodynamic and salinity variability normally caused by freshwater pulses. Monitoring data sets should be analyzed to quantify the drought's effect on the Delta. The Guadalquivir Estuary in Spain has a similar Mediterranean climate, and drought there increased salinity and displaced communities upstream (González–Ortegón et al. 2015).

Salinity management must consider fluvial, anthropogenic, and ocean drivers. Freshwater flow to the Delta depends primarily on precipitation, snowmelt, and runoff from the Central Valley watershed. Dam operations modify the outflow hydrograph, especially for low flows. In the Delta, gates, barriers, and exports modify flow paths and the salinity field. As sea level rises, ocean forcing in the form of potential salinity intrusion increases. This will require increased anthropogenic forcing to maintain the existing salinity field in the estuary. In other words, ocean forcing in the form of SLR will affect dam operations in the watershed and fluvial forcing on the Delta. Thus, the ocean, estuary, and watershed are linked with one another, and humans have created a feedback mechanism from the ocean to the watershed.

Fluvial and anthropogenic forcing have combined to greatly alter sediment supply to the Delta, and the magnitude of the change is similar to other estuaries. The pattern of human disturbance, increased sediment supply, dam construction, and decreased sediment supply is not uncommon for the Delta. Sediment supply to the Delta increased about an order of magnitude because of hydraulic mining in the late 1800s (Gilbert 1917). Sedimentation rates increased 2- to 10-fold in other California estuaries in the 19th and 20th centuries (Warrick and Farnsworth 2009). These increases are typical of the 5- to 10-fold increase found in lake and marine sediment records downstream from disturbed watersheds (Dearing and Jones 2003). Sediment supply in the Sacramento River decreased about 50% from 1957 to 2001 (Wright and Schoellhamer 2004). This magnitude of decrease is not uncommon; river sediment discharge to the coastal zone has decreased 45% in southern California from trapping behind dams (Warrick and Farnsworth 2009), 50% to 70% from the Mississippi River (Kesel 2003), 75% from the Trinity River in Texas (Ravens et al. 2009), and, globally, riverine sediment discharge to oceans has decreased  $10 \pm 2\%$ (Syvitski et al. 2005). Reforestation and dams have reduced the sediment discharge in the Changjiang (Yangtze River) 68% from the 1950s to 2000s, and the decrease is expected to reach 82% as dams

become fully operational and the system adjusts to them (Hu et al. 2009).

Several mechanisms are responsible for the increasing clarity of Delta waters (Hestir et al. 2013) and clearer waters affect the Delta ecosystem. Decreasing sediment supply from the watershed, floods that export sediment, and sedimentation in SAV beds increase water clarity. Increased water clarity, in turn, may contribute to increased harmful algal blooms (Jassby et al. 2003), SAV expansion (Santos et al. 2009, 2012), and decreased fish abundances (Sommer et al. 2007).

In addition to turbidity, endangered Delta Smelt are affected by water movement, temperature, and salinity. Recent physical advances of significance to understanding and managing Delta Smelt include the ability of backwater sloughs (isolated dead-end channels) to trap sediment and increase turbidity (Morgan–King and Schoellhamer 2013), the possible role of SAV in decreasing turbidity (Hestir et al. 2016), and the role of first flush and tidal currents on their landward migration (Bennett and Burau 2015). Though restoration efforts focus on marsh habitats, backwater sloughs may be an important and scarce habitat for Delta Smelt worthy of restoration.

Marsh restoration is an anthropogenic driver, and its success depends on physical factors such as fluvial sediment supply, a nondestructive wind wave environment, and optimal tidal inundation for plant colonization and growth. Tides, channel geometry, and marsh vegetation may interact to retain sediment in restoration sites. Increasing sediment retention in the Delta would decrease sediment supply downstream to the bay.

## **Information Gaps**

Here, we present what we believe are the most significant information gaps on flow and transport in the Delta, in roughly downstream order.

Though there is general understanding that sediment loads to the Delta have decreased over the past half century, likely from sediment trapping in reservoirs and other factors, the specific drivers of this trend are not understood. In particular, the current and historical watershed sources of sediment to the Delta, and how these sources have changed through time from anthropogenic influences in the watershed, are not understood. Because deposited sediment in the Delta essentially records the history of sediment sources, sediment "fingerprinting," whereby the geochemical signatures of source and sink sediments are examined and linked, could fill this information gap. Changes in watershed sources, such as the elimination of sources upstream from dams, should be reflected in the sediments deposited in the Delta over time, assuming the geochemical signatures of sediments from different regions of the watershed are significantly different. In addition, the currentday supply of sediment from the Feather–Yuba–Bear watersheds is a data gap that hinders definitive analysis of sediment sources.

Water residence time is a key ecological variable that varies tremendously within the Delta. Some channels efficiently transport water and its constituents to the bay or pumps, and some dead-end sloughs have much smaller tidal excursions and longer residence times. Residence time may be particularly significant for the fate and adverse effects of toxic substances. The pristine Delta contained not only much more marsh but many more small dead-end channels, and undoubtedly had regions of longer overall residence time than today's channelized Delta. Additional studies that examine residence time at the regional scale are needed to identify opportunities where increasing residence time improves habitat quality, and pelagic primary production.

To restore pelagic productivity in the Delta, the links between increasing water clarity, nutrient availability, residence time, and phytoplankton production need to be better understood (Cloern 2007; Lopez et al. 2006). Similarly, the relationship between flow characteristics such as velocity and turbulence and the production and sedimentation of organicrich particles (phytoplankton, phytodetritus, and detritus) has not been well characterized for Delta environments; a mismatch between production and transport may lead to significant attenuation of this carbon supply to lower Delta food webs (Lucas and Thompson 2012).

SAV has increased dramatically in the Delta, particularly in the south Delta, which appears to trap sediment and increase water clarity. SAV, floating vegetation, and decreased sediment supply from the watershed may explain observed decreases in turbidity (Hestir et al. 2016). Given the potential strong feedback between increased clarity and increased SAV, these changes may be an important aspect of the future Delta. Thus, studies are needed to understand the effects of SAV on Delta sediment transport and turbidity.

A question that may be difficult to answer definitively is whether the Delta is now more or less turbid than it was before the Gold Rush when Delta Smelt were presumably more abundant. Dams, deforestation, mining, urbanization, agriculturalization, and river channelization have all changed radically since the Gold Rush, and affect sediment supply to the Delta. Reshaping of the Delta landscape likely altered sediment transport. With no data from before the Gold Rush, numerical models of the watershed and Delta in the early 1800s are likely the best albeit speculative approach to answer this question.

Expanding the existing temperature sensor network to include near-bottom as well as near-surface temperatures would enable monitoring of temperature stratification and near-bottom cooler water that could serve as daytime refugia for fish.

Prevention of salinity intrusion during drought is currently a topic of great interest that also merits additional study to determine how to efficiently prevent intrusion while minimally disrupting the ecosystem. Each barrier placement should be considered an experiment from which analysis of data collected before and after placement and removal would provide lessons for building a better barrier next time.

Uncertainty in low flow values of Delta discharge likely plays a major role in defining the uncertainty in forecasting low flow values of X2. A significant effort to reduce the uncertainty in low-flow values of Delta discharge is needed.

# CONCLUSIONS

The tremendous expansion of acoustic and optical instruments deployed in the Delta over the past decade has greatly improved our understanding of how tidal variability affects flow and transport. Low Delta outflows are poorly estimated by a tidally

#### DECEMBER 2016

averaged water balance. Water quality sampling that fails to consider tidal variability can bias results. Transport in the Delta is driven more by tidal flow than tidally averaged flow. Mixing at junctions disperses water-quality constituents. Gates and barriers can significantly alter transport pathways. In addition, storms with similar time-scales of hours can supply and transport significant quantities of sediment and water-quality constituents.

Sediment is increasingly viewed as a diminishing resource needed to sustain the Delta ecosystem. Since 2000, the turbidity of Delta waters has decreased at an alarming rate. Native fish, including endangered Delta Smelt, favor turbid waters. Sediment supply is a key variable in determining the sustainability of tidal marsh as sea level rises.

Connections among the watershed, Delta, and the bay that have been quantified recently highlight that a landscape view of this system—rather than consideration of each region in isolation—is often needed. Reservoirs release water to maintain a salinity field in the Delta and bay that allows water diversions, and the volume of water needed to accomplish this will increase as sea level rises. The first flush of the watershed at the beginning of the wet season delivers a pulse of sediment and other constituents that affect the water quality and ecology of the Delta. The timing and magnitude of freshwater discharge from the Delta to the bay determines flushing and transport in south San Francisco Bay.

# ACKNOWLEDGEMENTS

We thank the Delta Science Program for their support and the Editors of the State of Bay–Delta Science 2016 for their guidance. Richard Norgaard, Michael Healey, Paul Work, and three anonymous reviewers provided constructive comments to previous manuscripts.

# REFERENCES

Alpers CN, Fleck JA, Marvin–DiPasquale M, Stricker CA, Stephenson M, Taylor HE. 2014. Mercury cycling in agricultural and managed wetlands, Yolo bypass, California: Spatial and seasonal variations in water quality. Sci Tot Environ 484(15):276–287. doi: http://dx.doi.org/10.1016/j.scitotenv.2013.10.096

- Anohin V, Imberger J, Romero J, Ivey G. 2006. Effect of long internal waves on the quality of water withdrawn from a stratified reservoir. J Hydraul Eng 132(11):1134–1145. doi: http://dx.doi.org/10.1061/ (ASCE)0733-9429(2006)132:11(1134)
- Bennett WA, Burau JR. 2015. Riders on the storm: selective tidal movements facilitate the spawning migration of threatened Delta Smelt in the San Francisco Estuary. Estuaries Coasts 38(3):826–835. doi: http://dx.doi.org/10.1007/s12237-014-9877-3
- Bergamaschi BA, Fleck JA, Downing BD, Boss E, Pellerin B, Ganju NK, Schoellhamer DH, Byington AA, Heim WA, Stephenson M, Fujii R. 2011. Methyl mercury dynamics in a tidal wetland quantified using in situ optical measurements. Limnol Oceanogr 56(4):1355– 1371. http://dx.doi.org/10.4319/lo.2011.56.4.1355
- Bergamaschi BA, Fleck JA, Downing BD, Boss E, Pellerin BA, Ganju NK, Schoellhamer DH, Byington AA, Heim WA, Stephenson M, Fujii R. 2012. Mercury dynamics in a San Francisco Estuary tidal wetland: assessing dynamics using in situ measurements. Estuaries Coasts 35(4):1036–1048.

doi: http://dx.doi.org/10.1007/s12237-012-9501-3

- Bergamaschi BA, Kuivila KM, Fram MS, 2001. Pesticides associated with suspended sediments entering San Francisco Bay following the first major storm of water year 1996. Estuaries 24(3):368–380. doi: http://dx.doi.org/10.2307/1353239
- Brennan ML, Schoellhamer DH, Burau JR, Monismith SG. 2002. Tidal asymmetry and variability of cohesive sediment transport at a site in San Francisco
  Bay, California. [accessed 2016 December 21]. In: Winterwerp JC, Kranenburg C, editors. INTERCOH–2000: fine sediment dynamics in the marine environment. Proceedings in Marine Science 5. Amsterdam u.a.: Elsevier p. 93–108. Available from: http://ca.water.usgs. gov/user\_projects/sfbay/publications\_group/brennan\_et\_al\_tidal.pdf
- Burau J, Ruhl C, Work P. 2016. Innovation in monitoring: the U.S. Geological Survey Sacramento–San Joaquin River Delta, California, flow station network. U.S. Geological Survey Fact Sheet 2015–3061. 6 p. doi: http://dx.doi.org/10.3133/fs20153061
- Chen SN. 2015. Asymmetric estuarine responses to changes in river forcing: a consequence of nonlinear salt flux. J Phys Oceanogr 45:2836–2847. doi: http://dx.doi.org/10.1175/JPO-D-15-0085.1

Cheng RT, Casulli V, Gartner JW. 1993. Tidal, residual, intertidal mudflat (TRIM) model and its applications to San Francisco Bay, California. Estuar Coastal Shelf Sci 36(3):235–280.

doi: http://dx.doi.org/10.1006/ecss.1993.1016

- Chua, VP, Xu M. 2014. Impacts of sea-level rise on estuarine circulation: an idealized estuary and San Francisco Bay. J Mar Syst 139:58–67. doi: http://dx.doi.org/10.1016/j.jmarsys.2014.05.012
- Cloern, JE. 2007. Habitat connectivity and ecosystem productivity: implications from a simple model. Am Nat 169(1):E21–33. Available from: *http://www.jstor.org/ stable/10.1086/510258*
- Cloern JE, Knowles N, Brown LR, Cayen D, Dettinger MD, Morgan TL, Schoellhamer DH, Stacey M, van der Wegen M, Wagner WR, Jassby AD. 2011. Projected evolution of California's San Francisco Bay–Delta River System in a century of climate change. PLoS ONE 6(9). doi: http://dx.doi.org/10.1371/journal.pone.0024465
- Dearing JA, Jones RT. 2003. Coupling temporal and spatial dimensions of global sediment flux through lake and marine sediment records. Glob Planet Change 39:147–168.

doi: http://dx.doi.org/10.1016/S0921-8181(03)00022-5

- Domagalski JL, Alpers CN, Slotton DG, Suchanek TH, Ayers SM. 2004. Mercury and methylmercury concentrations and loads in the Cache Creek watershed, California. Sci Total Environ 327(1–3):215–237. doi: http://dx.doi.org/10.1016/j.scitotenv.2004.01.013
- Downing BD, Boss E, Bergamaschi BA, Fleck JA, Lionberger MA, Ganju NK, Schoellhamer DH, Fujii R. 2009. Quantifying fluxes and characterizing compositional changes of dissolved organic matter in aquatic systems in situ using combined acoustic and optical measurements. Limnol Oceanogr Methods 7(1):119-131. doi: http://dx.doi.org/10.4319/ lom.2009.7.119
- Fischer H, List E, Koh R, Imberger J, Brooks N. 1979. Mixing in inland and coastal waters. San Diego (CA): Academic Press.
- Florsheim JL, Pellerin BA, Oh NH, Ohara, N, Bachand PAM, Bachand SM, Bergamaschi BA, Hernes PJ, Kavvas ML. 2011. From deposition to erosion: spatial and temporal variability of sediment sources, storage, and transport in a small agricultural watershed. Geomorphology 132(3–4):272–286. doi: http://dx.doi.org/10.1016/j. geomorph.2011.04.037

- Fong DA, Monismith SG, Stacey MT, Burau JR. 2009. Turbulent stresses and secondary currents in a tidal-forced channel with significant curvature and asymmetric Bed Forms. J Hydraul Eng 135(3):198-208. doi: http://dx.doi.org/10.1061/ (ASCE)0733-9429(2009)135:3(198)
- Ganju NK, Schoellhamer DH, Bergamaschi BA. 2005.
  Suspended sediment fluxes in a tidal wetland: measurement, controlling factors, and error analysis.
  Estuaries 28(6):812-822. doi: http://dx.doi.org/10.1007/ BF02696011
- Gilbert GK. 1917. Hydraulic mining debris in the Sierra Nevada. [Internet]. [accessed 2016 Dec 14]. U.S.
  Geological Survey Professional Paper 105. Available from: https://pubs.er.usgs.gov/publication/pp105
- Gleichauf KT. 2015. Temperature and tidal river junction dynamics in the Sacramento-San Joaquin Delta [PhD dissertation]. [Stanford (CA)]: Dept. of Civil and Environmental Engineering, Stanford University. 230 p. Available from: https://purl.stanford.edu/dg848xb0777
- Gleichauf, KT, Wolfram PJ, Monsen NE, Fringer OB, Monismith, SG. 2014. Dispersion mechanisms of a tidal river junction in the Sacramento-San Joaquin Della, California. San Franc Estuary Watershed Sci 12(4). doi: http://dx.doi.org/10.15447/sfews.2014v12iss4art1
- González-Ortegón E, Baldó F, Arias A, Cuesta JA, Fernández-Delgado C, Vilas C, Drake P. 2015. Freshwater scarcity effects on the aquatic macrofauna of a European Mediterranean-climate estuary. Sci Tot Environ 503-504:213-221.

doi: http://dx.doi.org/10.1016/j.scitotenv.2014.06.020

Grimaldo LF, Sommer T, Van Ark N, Jones G, Holland E, Moyle PB, Smith P. 2009. Factors affecting fish entrainment into massive water diversions in a tidal freshwater estuary: can fish losses be managed?. N Am J Fish Manage 29(5):1253-1270. doi: http://dx.doi.org/10.1577/M08-062.1

Healey MC, Dettinger MD, Norgaard RB, editors. 2008. The State of Bay–Delta Science 2008. [Internet]. [accessed 2016 Dec 14]. Sacramento (CA): CALFED Science Program. 174 p. Available from: *http://www.science. calwater.ca.gov/pdf/publications/sbds/sbds\_2008\_final\_ report\_101508.pdf*  Healey M, Goodwin P, Dettinger M, Norgaard R. 2016. The State of Bay-Delta Science 2016: an introduction. San Franc Estuary Watershed Sci 14(2).
doi: http://dx.doi.org/10.15447/sfews.2016v14iss2art5

Hestir EL, Schoellhamer DH, Greenberg J, Morgan TL, Ustin SL. 2016. The effect of submerged aquatic vegetation expansion on a declining turbidity trend in the Sacramento-San Joaquin River Delta. Estuaries Coasts 39(4):1100-1112. doi: http://dx.doi.org/10.1007/s12237-015-0055-z

Hestir EL, Schoellhamer DH, Morgan TL, Ustin SL. 2013. A step decrease in sediment concentration in a highly modified tidal river delta following the1983 El Niño floods. Mar Geol 345:304-313. doi: http://dx.doi.org/10.1016/j.margeo.2013.05.008

Holleman R, Stacey MT. 2014. Coupling of sea level rise, tidal amplification, and inundation. J Phys Ocean 44:1439-1455. doi: http://dx.doi.org/10.1175/JPO-D-13-0214.1

Hu B, Yang Z, Wang H, Sun X, Bi N, Li G. 2009.
Sedimentation in the Three Gorges Dam and the future trend of Changjiang (Yangtze River) sediment flux to the sea. Hydrol Earth Syst Sci 13: 2253–2264.
doi: http://dx.doi.org/10.5194/hess-13-2253-2009

Huisman J, van Oostveen P, Weissing F. 1999. Critical depth and critical turbulence: two different mechanisms for the development of phytoplankton blooms. Limnol Oceanogr 44(7):1781-1787.
doi: http://dx.doi.org/10.4319/lo.1999.44.7.1781

Jassby AD, Cloern JE, Mueller-Solger A. 2003. Phytoplankton fuels Delta food web. Cal Agric 57:104-109. doi: http://dx.doi.org/10.3733/ca.v057n04p104

Jassby A, Kimmerer W, Monismith S, Armor C, Cloern J, Powell T, Schubel J, Vendlinski T. 1995. Isohaline position as a habital indicator for estuarine populations. Ecol App 5:272-289. doi: http://dx.doi.org/10.2307/1942069

Jones NL, Monismith SG. 2008. The influence of whitecapping waves on the vertical structure of turbulence in a shallow estuarine embayment. J Phys Oceanogr 38(7): 1563–1580. doi: http://dx.doi.org/10.1175/2007JP03766.1 Jones NL, Thompson JK, Monismith SG. 2008. A note on the effect of wind waves on vertical mixing in Franks Tract, Sacramento-San Joaquin Delta, California. San Franc Estuary Watershed Sci 6(2). doi: http://dx.doi.org/10.15447/sfews.2008v6iss2art4

Jones NL, Thompson JK, Arrigo KR, Monismith SG. 2009. Hydrodynamic control of phytoplankton loss to the benthos in an estuarine environment. Limnol Oceanogr 54(3):952-969.

doi: http://dx.doi.org/10.4319/lo.2009.54.3.0952

Kesel RH. 2003. Human modifications to the sediment regime of the Lower Mississippi river flood plain. Geomorphology 56:325–334. doi: http://dx.doi.org/10.1016/S0169-555X(03)00159-4

Kimmerer W. 2008. Losses of Sacramento River Chinook Salmon and Delta Smelt to entrainment in water diversions in the Sacramento-San Joaquin Delta. San Franc Estuary Watershed Sci 6(2). doi: http://dx.doi.org/10.15447/sfews.2008v6iss2art2

Kimmerer WJ, MacWilliams ML, Gross ES. 2013. Variation of fish habitat and extent of the low-salinity zone with freshwater flow in the San Francisco Estuary. San Franc Estuary Watershed Sci 11(4). doi: http://dx.doi.org/10.15447/sfews.2013v11iss4art1

Kimmerer W, Nobriga M. 2008. Investigating particle transport and fate in the Sacramento-San Joaquin Delta using a particle tracking model. San Franc Estuary Watershed Sci 6(1).

doi: http://dx.doi.org/10.15447/sfews.2008v6iss1art4

Lehman PW, Mayr S, Mecum L, Enright C. 2009. The freshwater tidal wetland Liberty Island, CA, was both a source and sink of inorganic and organic material to the San Francisco Estuary. Aquat Ecol 44(2):359-372. doi: http://dx.doi.org/10.1007/s10452-009-9295-y

Lopez CB, Cloern JE, Schraga TS, Little AJ, Lucas LV, Thompson JK, Burau JR. 2006. Ecological values of shallow-water habitats: Implications for the restoration of disturbed ecosystems. Ecosystems 9(3):422-440. doi: http://dx.doi.org/10.1007/s10021-005-0113-7

Lucas LV, Sereno DM, Burau, JR, Schraga TS, Lopez CB, Stacey MT, Parchevsky KV, Parchevsky VP. 2006. Intradaily variability of water quality in a shallow tidal lagoon: mechanisms and implications. Estuaries Coasts 29(5):711-730.

doi: http://dx.doi.org/10.1007/BF02786523

- Lucas LV, Thompson JK. 2012. Changing restoration rules: Exotic bivalves interact with residence time and depth to control phytoplankton productivity. Ecosphere 3(12). doi: http://dx.doi.org/10.1890/ES12-00251.1
- MacIntyre S. 1993. Vertical mixing in a shallow, eutrophic lake: Possible consequences for the light climate of phytoplankton. Limnol Oceanogr 38:798–817. doi: http://dx.doi.org/10.4319/lo.1993.38.4.0798
- Marineau MD, Wright SA. Forthcoming. Bed-Material Characteristics of the Sacramento-San Joaquin River Delta, California, 2010-13. USGS Data Series Report.
- McCulloch DS, Peterson DH, Carlson PR, Conomos TJ. 1970. A preliminary study of the effects of water circulation in the San Francisco Bay Estuary: some effects of fresh-water inflow on the flushing of South San Francisco Bay. [accessed 2016 December 21]. U.S. Geological Survey Circular 637A. 27 p. Available from: https://pubs.er.usgs.gov/publication/cir637AB
- McKee LJ, Lewicki M, Schoellhamer DH, Ganju NK. 2013. Comparison of sediment supply to San Francisco Bay from coastal and Sierra Nevada watersheds. Mar Geol 345:47-62.

doi: http://dx.doi.org/10.1016/j.margeo.2013.03.003

- Moftakhari HR, Jay DA, Talke SA, Kukulka T, Bromirski PD. 2013. A novel approach to flow estimation in tidal rivers. Water Resour Res 49(8):4817– 4832. doi: http://dx.doi.org/10.1002/wrcr.20363
- Moftakhari HR, Jay DA, Talke SA, Schoellhamer DH. 2015. Estimation of historic flows and sediment loads to San Francisco Bay, 1849-2011. J Hydrol 529(3):1247-1261. doi: http://dx.doi.org/10.1016/j.jhydrol.2015.08.043
- Monismith SG. Forthcoming. An integral model of unsteady salinity intrusion in estuaries. *Journal of Hydraulic Research*.
- Monismith SG, Hench JL, Fong DA, Nidzieko NJ, Fleenor WE, Doyle L, Schladow SG. 2009. Thermal variability in a tidal river. Estuaries Coasts 32:100–110. doi: http://dx.doi.org/10.1007/s12237-008-9109-9

- Monismith S, Hench J, Smith P, Fleenor W, Doyle L, Schladow SG. 2008. An application of the SI3D hydrodynamics model to the Stockton Deep Water Ship Channel: physics and model application [accessed 2016 December 21]. In: Schladow SG, Monismith S, editors. Hydrodynamics and oxygen modeling of the Stockton Deep Water Ship Channel, Final report to CALFED. Project ERP-02D-P51. p. 74-133. Available from: https:// watershed.ucdavis.edu/pdf/Schladow-Monismith-2009. pdf
- Monismith SG, Kimmerer W, Stacey MT, Burau JR. 2002. Structure and flow-induced variability of the subtidal salinity field in northern San Francisco Bay. J Phys Ocean 32(11):3003-3019. doi: http://dx.doi. org/10.1175/1520-0485(2002)032%3C3003:SAFIV0% 3E2.0.C0;2
- Monsen N. 2000. A study of sub-tidal transport in Suisun Bay and the Sacramento-San Joaquin Delta, California. [PhD dissertation]. [Stanford (CA)]: Dept. of Civil and Environmental Engineering, Stanford University.
- Monsen NE, Cloern JE, Burau JR. 2007. Effects of flow diversions on water and habitat quality: examples from California's highly manipulated Sacramento-San Joaquin Delta. San Franc Estuary Watershed Sci 5(3). doi: http://dx.doi.org/10.15447/sfews.2007v5iss5art2
- Morgan–King TL, Schoellhamer DH. 2013. Suspendedsediment flux and retention in a backwater tidal slough complex near the landward boundary of an estuary. Estuaries Coasts 36(2):300–318. doi: http://dx.doi.org/10.1007/s12237-012-9574-z
- Morgan–King T, Wright S. 2015. Sediment budgets, transport, and depositional trends in a large tidal delta. Proceedings of the 10th Federal Interagency Sedimentation Conference; 2015 April 19–22; Reno, NV.
- Moskalski SM, Torres R, Bizimis M, Goni M, Bergamaschi B, Fleck J. 2013. Low-tide rainfall effects on metal content of suspended sediment in the Sacramento-San Joaquin Delta. Cont Shelf Res 56:39-55. doi: http://dx.doi.org/10.1016/j.csr.2013.02.001
- Moyle PB, Brown LR, Durand JR, Hobbs JA. 2016. Delta Smelt: life history and decline of a once abundant species in the San Francisco Estuary. San Franc Estuary Watershed Sci 14(2).

doi: http://dx.doi.org/10.15447/sfews.2016v14iss2art6

Nobriga ML, Feyrer F, Baxter RD, Chotkowski M. 2005. Fish community ecology in an altered river delta: spatial patterns in species composition, life history strategies, and biomass. Estuaries 5:776–785. doi: http://dx.doi.org/10.1007/BF02732915

O'Brien KR, Meyer DL, Waite AM, Ivey GN, Hamilton DP. 2004. Disaggregation of *Microcystis aeruginosa* colonies under turbulent mixing: laboratory experiments in a grid-stirred tank. Hydrobiologica 519:143–152. doi: *http://dx.doi.org/10.1023/ B:HYDR.0000026501.02125.cf* 

- Oltmann R. 1995. Continuous flow measurements using ultrasonic velocity meters: an update. IEP Newsletter 8(4):22-25. [Internet]. [accessed 2016 Dec 14]. Available from: http://www.water.ca.gov/iep/newsletters/1995/fall/ page22.pdf
- Oltmann RN. 1998. Indirect measurement of delta outflow using ultrasonic velocity meters and comparison with mass-balance calculated outflow. IEP Newsletter 11(1). [Internet]. [accessed 2016 Dec 14]. Available from: http://www.water.ca.gov/iep/newsletters/1998/ winter/Indirect%20Measurement%20of%20Delta%20 Outflow%20Using%20Ultrasonic%20Velocity%20 Meters.pdf
- Oh NH, Pellerin BA, Bachand PAM, Hernes PJ, Bachand SM, Ohara N, Kavvas ML, Bergamaschi BA, Horwath WR. 2013. The role of irrigation runoff and winter rainfall on dissolved organic carbon loads in an agricultural watershed. Agric Ecosyst Environ 179:1-10. doi: http://dx.doi.org/10.1016/j.agee.2013.07.004
- Paerl H., Huisman J. 2009. Climate change: a catalyst for global expansion of harmful cyanobacterial blooms. Env Micro Reports 1(1):27–37. doi: http://dx.doi. org/10.1111/j.1758-2229.2008.00004.x
- Pellerin BA, Bergamaschi BA, Downing BD, Saraceno JF, Garrett JD, Olsen LD. 2013. Optical techniques for the determination of nitrate in environmental waters: guidelines for instrument selection, operation, deployment, maintenance, quality assurance, and data reporting. [Internet]. [accessed 2016 Dec 14]. U.S. Geological Survey Techniques and Methods 1-D5. Available from: http://pubs.usgs.gov/tm/01/d5/

- Pellerin BA, Downing BD, Kendall C, Dahlgren RA, Kraus TEC, Saraceno J, Spencer RGM, Bergamaschi BA. 2009. Assessing the sources and magnitude of diurnal nitrate variability in the San Joaquin River (California) with an *in situ* optical nitrate sensor and dual nitrate isotopes. Freshw Biol 54(2):376-387. doi: *http://dx.doi. org/10.1111/j.1365-2427.2008.02111.x*
- Perry RW, Buchanan RA, Brandes PL, Burau JR, Israel JA. 2016. Anadromous salmonids in the Delta: new science 2006–2016. San Franc Estuary Watershed Science 14(2). doi: http://dx.doi.org/10.15447/sfews.2016v14iss2art7
- Ravens TM, Thomas RC, Roberts KA, Santshi PH. 2009. Causes of salt marsh erosion in Galveston Bay, Texas. J Coast Res 25:265–272. doi: http://dx.doi.org/10.2112/07-0942.1
- Ridderinkhof H, Zimmerman JTF. 1992. Chaotic stirring in a tidal system. Science 258 (5085):1107-1109. doi: http://dx.doi.org/10.1126/science.258.5085.1107
- Robarts RD, Zohary T. 1987. Temperature effects on photosynthetic capacity, respiration, and growth rates of bloom-forming cyanobacteria. NZ J Mar Freshw Res 21(3):391-399.

doi: http://dx.doi.org/10.1080/00288330.1987.9516235

- Roth DA, Taylor HE, Domagalski J, Dileanis P, Peart DB, Antweiler RC, Alpers CN. 2001. Distribution of inorganic mercury in Sacramento River water and suspended colloidal sediment material. Arch Environ Contam Toxicol 40(2):161-172. doi: http://dx.doi.org/10.1007/s002440010159
- Ruhl CA, Simpson MR. 2005. Computation of discharge using the index-velocity method in tidally affected areas. [Internet]. [accessed 2016 Dec 14]. U.S. Geological Survey Scientific Investigations Report 2005-5004. Available from: http://pubs.er.usgs.gov/publication/ sir20055004
- Russo TA, Fisher AT, Winslow DM. 2013. Regional and local increases in storm intensity in the San Francisco Bay Area, USA, between 1890 and 2010. J Geophys Res Atmos 118:3392-3401. doi: http://dx.doi.org/10.1002/ jgrd.50225
- Santos MJ, Hestir EL, Khanna S, Ustin SL. 2012. Image spectroscopy and stable isotopes elucidate functional dissimilarity between native and nonnative plant species in the aquatic environment. New Phytol 193:683-695. doi: http://dx.doi. org/10.1111/j.1469-8137.2011.03955.x

Santos MJ, Khanna S, Hestir EL, Andrew ME, Rajapakse S, Greenberg JA, Anderson LW, Ustin SL. 2009. Use of hyperspectral remote sensing to evaluate efficacy of aquatic plant management. Invasive Plant Sci Manag 2:216-229.

doi: http://dx.doi.org/10.1614/IPSM-08-115.1

Saraceno JF, Pellerin, BA, Downing, BD, Boss, E, Bachand PAM, Bergamaschi BA. 2009. High-frequency *in situ* optical measurements during a storm event: Assessing relationships between dissolved organic matter, sediment concentrations, and hydrologic processes. J Geophys Res 114. doi: *http://dx.doi.org/10.1029/2009JG000989* 

Schoellhamer DH, Wright SA, Drexler JZ. 2012. Conceptual model of sedimentation in the Sacramento–San Joaquin River Delta. San Franc Estuary Watershed Sci 10(3). doi: http://dx.doi.org/10.15447/sfews.2012v10iss3art3

Schoellhamer DH, Wright SA, Drexler JZ. 2013. Adjustment of the San Francisco estuary and watershed to reducing sediment supply in the 20th century. Mar Geol 345:63-71. doi: http://dx.doi.org/10.1016/j.margeo.2013.04.007

Shellenbarger GG, Wright SA, Schoellhamer DH. 2013. A sediment budget for the southern reach in San Francisco Bay, CA: implications for habitat restoration.
Mar Geol 345:281-293.
doi: http://dx.doi.org/10.1016/j.margeo.2013.05.007

Simpson MR, Oltmann RN. 1993. Discharge-measurement system using an acoustic Doppler current profiler with applications to large rivers and estuaries. [Internet]. [accessed 2016 Dec 14]. U.S. Geological Survey Water-Supply Paper 2395. 32 p. Available from: http://pubs. er.usgs.gov/publication/wsp2395

Singer MB. 2007. The influence of major dams on hydrology through the drainage network of the Sacramento River basin, California. River Res Appl 23:55–72. doi: *http://dx.doi.org/10.1002/rra.968* 

Sommer TR, Armor C, Baxter RD, Breuer R, Brown L, Chotkowski M, Culberson S, Feyrer F, Gingras M, Herbold B, Kimmerer W, Mueller-Solger A, Nobriga ML, Souza K. 2007. The collapse of pelagic fishes in the upper San Francisco estuary. Fisheries 32:270-277. doi: http://dx.doi.org/10.1577/1548-8446(2007)32[270:TC0 PFI]2.0.C0;2 Sridharan VK, 2015, Scalar transport in channel networks: development of a particle tracking model to study the movement of scalars in the Sacramento-San Joaquin Delta [PhD dissertation]. [Stanford (CA)]: Dept. of Civil and Environmental Engineering, Stanford University.
476 p. Available from: https://purl.stanford.edu/ kp594nf8504

Swanson KM, Drexler JZ, Fuller C, Schoellhamer DH. 2015. Modeling tidal freshwater marsh sustainability in the Sacramento-San Joaquin Delta under a broad suite of potential future scenarios. San Franc Estuary Watershed Sci 13(1).

doi: http://dx.doi.org/10.15447/sfews.2015v13iss1art3

Syvitski JPM, Vorosmarty CJ, Kettner AJ, Green P. 2005. Impact of humans on the flux of terrestrial sediment to the global coastal ocean. Science 308:376–380. doi: http://dx.doi.org/10.1126/science.1109454

Wagner RW, Stacey MT, Brown LR, Dettinger M. 2011. Statistical models of temperature in the Sacramento-San Joaquin Delta under climate-change scenarios and ecological implications. Estuaries Coasts 34:544-556. doi: http://dx.doi.org/10.1007/s12237-010-9369-z

Wang J, Yin H, Chung F. 2011. Isolated and integrated effects of sea level rise, seasonal runoff shifts, and annual runoff volume on California's largest water supply. J Hydrology 405:83-92. doi: http://dx.doi.org/10.1016/j.jhydrol.2011.05.012

Warrick JA, Farnsworth KL. 2009. Sources of sediment to the coastal waters of the southern California bight. In: Earth science in the urban ocean: the southern California borderland. Lee HJ, Normark WR, editors. Geolog Soc Am Spec Paper 454:39–52. doi: http://dx.doi.org/10.1130/2009.2454(2.2)

Weston DP, Asbell AM, Lesmeister SA, Teh SJ, Lydy MJ. 2014. Urban and agricultural pesticide inputs to a critical habitat for the threatened delta smelt (*Hypomesus transpacificus*). Environ Toxicol Chem 33(4):920-929. doi: http://dx.doi.org/10.1002/etc.2512

Weston DP, You J, Amweg EL, Lydy MJ. 2008. Sediment toxicity in agricultural areas of California and the role of hydrophobic pesticides. In: Gan J, Spurlock F, Hendley P, Weston DP, editors. Synthetic Pyrethroids 991:26-54. doi: http://dx.doi.org/10.1021/bk-2008-0991.ch002 Whipple AA, Grossinger RM, Rankin D, Stanford B, Askevold RA. 2012. Sacramento-San Joaquin Delta historical ecology Investigation: exploring pattern and process. [Internet]. [accessed 2016 Dec 14]. A report of SFEI-ASC's Historical Ecology Program, Publication #672. Richmond (CA): San Francisco Estuary Institute-Aquatic Science Center. Available from: http://www.sfei. org/documents/sacramento-san-joaquin-delta-historicalecology-investigation-exploring-pattern-and-proces

Wolfram PJ, Fringer OB, Monsen NE, Gleichauf KT, Fong DA, Monismith, SG. 2016. Modeling intrajunction dispersion at a well-mixed tidal river junction. J Hydraul Eng 142(8). doi: http://dx.doi.org/10.1061/(ASCE) HY.1943-7900.0001108

Wright SA, Morgan TL. 2015. Suspended sediment transport through a large fluvial-tidal channel network. Proceedings of the 10th Federal Interagency Sedimentation Conference; 2015 April 19-22; Reno, NV.

Wright SA, Schoellhamer DH. 2005. Estimating sediment budgets at the interface between rivers and estuaries with application to the Sacramento–San Joaquin River Delta. Water Resour Res 41. doi: http://dx.doi.org/10.1029/2004WR003753

# **NOTES**

List E. 1995. Personal communication with S. G. Monismith on dispersion associated with junctions in the Delta.