

Yolo Bypass Invertebrate Drift Sample Collection Meta Data – Lynn Takata

Suggested Citation: California Department of Water Resources, Aquatic Ecology Section. Yolo Bypass Invertebrate Drift Sample Collection Meta Data. 2016.

IEP Aquatic Ecology Section Yolo Bypass Invertebrate Drift Sample Collection Meta Data Last updated: August 2016

I. Contact Information

Program Manager: Brian Schreier

Lead Contact: Jared Frantzich
Dept. of Water Resources
Division of Environmental Services
3500 Industrial Blvd., West Sacramento, CA.
916-376-9823
email: jfrantzi@water.ca.gov

II. Study Elements and Objectives

Largely supported by the Interagency Ecological Program (IEP), DWR has operated a fisheries and invertebrate monitoring program in the Yolo Bypass since 1998. The project has provided a wealth of information regarding the significance of seasonal floodplain habitat to native fishes. Basic objectives of the project are to collect baseline data on lower trophic levels (phytoplankton, zooplankton and invertebrate drift), juvenile and adult fish, hydrology, and physical conditions. As the Yolo Bypass has been identified as a high restoration priority by the US Fish and Wildlife Service and National Marine Fisheries Service biological opinions for Delta Smelt (*Hypomesus transpacificus*) and winter and spring-run Chinook Salmon (*Oncorhynchus tshawytscha*), and by California EcoRestore, these baseline data are critical for evaluating success of future restoration projects. In addition, the data have already served to increase our understanding of the role of the Yolo Bypass in the life history of native fishes and its ecological function in the San Francisco Estuary. Key findings include: (1) Yolo Bypass is a major factor regulating year class strength of splittail, *Pogonichthys macrolepidotus* (Sommer et al., 1997; Feyrer et al., 2006; Sommer et al., 2007a); (2) Yolo Bypass is a key migration corridor for adult fish of several listed and sport fish (Harrell and Sommer 2003); (3) it is one of the most important regional rearing areas for juvenile Chinook Salmon (Sommer et al., 2001a; 2005); and (4) Yolo Bypass is a source of phytoplankton to the food web of the San Francisco Estuary (Jassby and Cloern 2000; Schemel et al., 2004; Sommer et al., 2004a).

The collection of invertebrate drift is one element of the Aquatic Ecology Section's (AES), Yolo Bypass Fish Monitoring Program's (YBFMP) lower trophic monitoring that is conducted under the IEP umbrella. The monitoring of invertebrate drift was initiated to compare the seasonal variations in densities and species trends of aquatic and terrestrial insects/non-insects within (1) Sacramento River channel, and (2) the Yolo Bypass, the river's seasonal floodplain. Aquatic and terrestrial insects are an important component in the diet of juvenile and adult fishes within the San Francisco Estuary, including two important native fishes: juvenile Chinook Salmon and Sacramento Splittail.

Key findings to date: (1) Chinook Salmon sampled in the floodplain had diets comprised of 90% Dipterans and zooplankton, with Chironomidae being the dominant Diptera family (Sommer et al., 2001), (2) The floodplain of the Yolo Bypass contains significantly higher densities of Diptera (Diptera densities being positively associated with flow) and terrestrial invertebrates than the adjacent Sacramento River (Sommer et al. 2001b; Sommer et al. 2004; Sommer et al. 2007), (3) A major portion of the diet of juvenile Sacramento Splittail are chironomid larvae (Kurth and Nobriga 2001, Moyle et al. 2004, Sommer et al. 2007), and (4) The Yolo Bypass was the site of the recent discovery of a new aestivating and winter emerging chironomid; *Hydrobaenus saetheri* (Cranston et al. 2007).

III. Study Area and Sample Sites


A. General Information

There are two fixed sampling site locations for this study: (1) Toe Drain of Yolo Bypass (STTD) at our rotary screw trap, and (2) Sacramento River at Sherwood Harbor (SHR). These sites are sampled on an ebb tide on the same day or within one day of one another.

B. Name and Location Information of Current Invertebrate Drift Sampling Sites

Station	Location	latitude			longitude			Start Year
		degrees	minutes	seconds	degrees	minutes	seconds	
STTD	Yolo Bypass - Screw Trap at Toe Drain	38	21	12.46	121	38	34.71	1998
SHR	Sacramento River at Sherwood Harbor	38	31	56.77	121	31	41.1	1998

Map of Currently Sampled Sites



IV. Period of Record

Invertebrate drift monitoring began in 1998 and continues through the present. The drift invertebrate dataset includes the proper sorting, identification, and enumeration of (1) aquatic insects, (2) aquatic non-insects, (3) terrestrial insects, and (4) terrestrial non-insects.

V. Sampling Frequency

Early in the monitoring program, sampling was generally conducted once monthly from February-April. Beginning in 2001, sampling was conducted at least once monthly during January-June. In some years, sampling was conducted weekly during the inundation and draining of the Yolo Bypass floodplain. Since 2011, sampling is conducted biweekly (every other week) year-round and weekly during floodplain inundation and drainage events.

Sampling Frequency by Month and Year

Yolo Bypass Screw Trap at Toe Drain (STTD)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
1998	0*	5*	4*	2*	0*	0*	0	0	0	0	0	0*	11
1999	0	2*	5*	3	0	0	0	0	0	0	0	0	10
2000	0	4*	3*	0	0	0	0	0	0	0	0	0	7
2001	1	2	2	2	1	0	0	0	0	0	0	0	8
2002	4*	2	2	2	3	2	0	0	0	0	0	0*	15
2003	6*	0	2	2	3*	2	0	0	0	0	0	0	15
2004	2*	2*	1*	2	1	1	0	0	0	0	0	1	10
2005	5	3	4	4	2*	2	0	0	0	0	0	0*	20
2006	3*	2*	4*	2*	2*	2	0	0	0	0	0	0	15
2007	0	1	2	2	2	2	0	0	0	0	0	0	9
2008	2	2	2	2	2	2	0	0	0	0	0	0*	12
2009	2	2	2	2	2	2	0	0	0	0	0	0	12
2010	4*	3	3	2	2	0	0	0	0	0	0	0*	14
2011	2*	2	2*	3*	2	3	2	2	2	2	3	2	27
2012	2	2	2	2	3	2	2	2	3	3	2	2*	27
2013	5	4	3	3	2	2	2	2	2	3	1	2	31
2014	3	2	2	2	2	2	2	3	2	2	2	3	27
2015	4	4	7	5	3	6	4	4	2	3	2	2	46
Total	46	44	52	42	32	30	12	13	11	13	10	12	316

*Months with overtopping at Fremont Weir.

Sacramento River at Sherwood Harbor (SHR)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
1998	0*	5*	4*	2*	1*	0*	0	0	0	0	0	0*	12
1999	0	2*	5*	3	0	0	0	0	0	0	0	0	10
2000	0	3*	3*	0	0	0	0	0	0	0	0	0	6
2001	1	2	2	2	1	0	0	0	0	0	0	0	8
2002	4*	2	2	2	3	2	0	0	0	0	0	0*	15
2003	6*	1	2	2	3	2	0	0	0	0	0	0	16
2004	1*	2*	1*	2	1	1	0	0	0	0	0	1	9
2005	5	3	4	4	2*	2	0	0	0	0	0	0*	20
2006	2*	2*	4*	2*	2*	2	0	0	0	0	0	0	14
2007	0	1	2	2	2	2	0	0	0	0	0	0	9
2008	2	2	2	2	2	1	0	0	0	0	0	0*	11
2009	2	2	2	2	2	2	0	0	0	0	0	0	12
2010	4*	4	3	2	2	0	0	0	0	0	0	0*	15
2011	2*	2	2*	3*	2	3	2	2	2	2	3	2	27
2012	2	2	2	2	3	2	2	2	2	3	2	2*	26
2013	5	4	3	3	2	2	2	2	2	3	1	2	31
2014	3	2	2	2	2	2	2	3	2	2	2	3	27
2015	4	4	6	5	4	4	5	5	2	2	2	2	45
Total	43	45	51	42	34	27	13	14	10	12	10	12	313

*Months with overtopping at Fremont Weir

Number of Sampling Events by Station and by Year (All Conducted Sampling Events)

Station	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Total
STTD	11	10	7	8	15	15	10	20	15	9	12	12	14	27	27	31	27	46	316
SHR	12	10	6	8	15	16	9	20	14	9	11	12	15	27	26	31	27	45	313
Total	23	20	13	16	30	31	19	40	29	18	23	24	29	54	53	62	54	101	629

VI. Field Collection Methods

An aquatic drift net is used to capture: (1) Aquatic insects (*Ephemeroptera*, *Trichoptera*, *Odonata*, *Hemiptera*, *Coleoptera*, and *Diptera*), (2) Aquatic non-insects (*Oligochaeta*, *Gastropoda*, *Amphipoda*, and *Ostracoda*), (3) Terrestrial insects (*Hemiptera*, *Coleoptera*, *Diptera*, and *Lepidoptera*) and (4) Terrestrial non-insects (*Mollusca*, *Acari* etc.).

Water quality parameters are recorded when the sample is collected. Temperature (C), electrical conductivity (uS/cm), dissolved oxygen (mg/L), and pH are measured using a YSI 556 Multiprobe System. Turbidity is measured from a water sample collected in a glass vial and later analyzed at the office using a Hach 2100Q Portable Turbidimeter. Secchi depth (cm) is also measured. Other factors including tide stage, weather, and trap condition code are also recorded.

A. Aquatic Drift Net

The aquatic drift net is made of 500 micron mesh net, with a 0.46 m by 0.3 m rectangular mouth and 0.91 m long, harnessed to a floated stainless steel frame. It tapers to 0.076 m at the cod-end where a polyethylene jar screened with 500 micron mesh collects the organisms. When there is sufficient flow (typically from January – June), Toe Drain samples are collected during the ebb tide from the rotary screw trap anchored in the middle of the channel and Sacramento River/Sherwood Harbor samples are taken dockside. In the absence of sufficient downstream flow, typically from July-Nov, Sacramento River and Yolo Bypass samples are taken from a boat moving approximately 2-3 mph upstream near the screw trap or dock. Net tow times have varied through the years, with shorter tows occurring with high flows or debris loads. Generally, tows have been 10 minutes long. Exact tow times are recorded with every sampling event.

The flow is measured with a General Oceanics Model 2030R flow meter. Samples are preserved in the field with 10% formalin with Rose Bengal dye to aid in separating organisms from detritus and algae.

VII. Lab Processing Methods

Current Procedure (1998-Current): All aquatic and terrestrial drift invertebrate samples are rinsed and passed through a 250 micron mesh sieve. Large debris (leaves, sticks, etc.) are carefully rinsed and removed with all the remaining material within the sieve being retained for identification. Within 2-3 weeks of collection, each invertebrate drift sample preserved in formalin in the field is transferred to 70-80% ETOH in the laboratory, for sorting, identification, and enumeration by the contractor: EcoAnalysts, Inc. (1420 South Blaine Street, Suite 14 Moscow, Idaho 83843). All the aquatic insects and non-insects are counted and identified to the family level. The terrestrial insects and non-insects are counted and identified to the order level.

Aquatic Drift Taxonomic Level Identification Table

AQUATIC INSECT		TERRESTRIAL NON-INSECT	
Ephemeroptera	Baetidae	Other Organisms	Araneae
Trichoptera	Hydroptilidae		Diplopoda
Odonata	Coenagrionidae		Geophilomorpha
	Gomphidae		
Hemiptera	Corixidae		
	Notonectidae		
AQUATIC NON-INSECT			
Annelida-Oligochaeta	Enchytraeidae		
	Naididae		
Mollusca-Gastropoda	Hydrobiidae		
	Lymnaeidae		
	Physidae		
	Planorbidae		
Crustacea-Amphipoda	Corophiidae		
	Crangonyctidae		
	Gammaridae		
	Hyalellidae		
Crustacea-Ostracoda	Ostracoda		
Acari	Acari		
	Arrenuridae		
	Limnesiidae		
	Oribatei		
Cnidaria	Hydridae		
Other Organisms	Nematoda		
TERRESTRIAL INSECT			
Hemiptera	Hemiptera		
Coleoptera	Coleoptera		
Diptera	Diptera		
Lepidoptera	Lepidoptera		
Other Insecta	Collembola		
	Hymenoptera		
	Neuroptera		
	Psocoptera		
	Thysanoptera		

The number per cubic meter for each aquatic and terrestrial organism taken in the aquatic drift net was calculated using the following equation:

$$N = C/V$$

Where:

N = the number of a taxon per cubic meter of water sampled

C = the total number of a taxon counted for the sample

V = the volume of water sampled through the net (m³)

Calculations for volume of water sampled through the net is specific to the General Oceanics Flowmeter model 2030R, and is calculated as follows (General Oceanics Inc.):

$$\frac{(\text{Flowmeter count start} - \text{Flowmeter count end}) \times \text{Rotor Constant}}{999999} \times \frac{\text{Net mouth area}}{4}$$

The rotor constant depends upon which the flowmeter rotor was used during each sampling event, and is identified in the sampling database. Rotor constants are specified in the General Oceanics Flowmeter 2030R manual as:

Standard Speed Rotor Constant = 26,873

Low Speed Rotor Constant R6 = 57,560

X. Data Management and Quality Assurance/Quality Control

A. Field Data

Field data are collected and recorded onto datasheets by DWR personnel. These data are then entered monthly by DWR personnel into an Access database. Field data are reviewed monthly for accuracy and completeness. Annually, after all samples are processed by the contractor for the year, lab data are reviewed for accuracy and completeness.

B. Field Datasheet

Paper datasheets are digitized and archived in binders that are stored at the West Sacramento, Industrial Blvd. DWR office.

Field Datasheet

LOWER TROPHIC SAMPLING – YOLO BYPASS STUDY

2015/2016

Location: _____

Crew: _____

Date: _____

Time: _____

pH: _____

DO: _____

SpCnd: _____

Cnd (EC): _____

Secchi Depth: _____ m

Water Temp: _____ °C

Weather: _____

Turb: _____ NTU

Vial #: _____

Light

Attenuation:

Surface Irradiance (in air avg): _____ μmol

Subsurface Irradiance (in water avg)(~75%, ~50%, ~25%, ~1%):

LICOR Calibration
-143.27 (in air)
-232.10 (in water)

0.75 = _____ μmol
0.50 = _____ μmol
0.25 = _____ μmol
0.01 = _____ μmol

Depth: _____ m

① _____ μmol

Depth: _____ m

② _____ μmol

Depth: _____ m

③ _____ μmol

Depth: _____ m

④ _____ μmol

Drift Sample :

Condition Code: _____

Start Time: _____

Stop Time: _____

Set Time: _____ min

Flow Meter:

Regular or Low Speed

Flow :

*For low speed, record initial meter reading in "end meter" box

Start Meter: _____

End Meter: _____

Comments: _____

Egg & Larval Fish Sample :

Condition Code: _____

1st Start Time: _____

1st Stop Time: _____

2nd Start Time: _____

2nd Stop Time: _____

Set Time: _____ min

Flow Meter:

Regular or Low Speed

*For low speed, record initial meter reading in "end meter" box

Flow :

(Mid-West) Start Meter: _____

End Meter: _____

(Near-West) Start Meter: _____

End Meter: _____

(Mid-East) Start Meter: _____

End Meter: _____

(Near-East) Start Meter: _____

End Meter: _____

Zooplankton Sample :

Condition Code: _____

Start Time: _____

150 Stop Time: _____

50 Stop Time: _____

150 Set Time: _____ min

50 Set Time: _____ min

Flow Meter:

Regular or Low Speed

*For low speed, record initial meter reading in "end meter" box

Flow :

150 μm : Start Meter: _____

End Meter: _____

50 μm : Start Meter: _____

End Meter: _____

Comments: _____

Chlorophyll Sample :

Yes ☐ No ☐

Replicate ☐

Time: _____

Filtered: 500mL OR 250mL

Phytoplankton Sample :

Yes ☐ No ☐

60 mL Amber Bottle w/ Lugol Solution

Entered by : _____

Date: _____

Checked by : _____

Date: _____

C. Taxonomic Data

Taxonomic results are received via email from the contractor, and entered into the AES Access database by DWR personnel. Electronic copies of results for taxonomic analyses are archived on DWR/AES Network drives. Hard copies are printed and archived in binders at the West Sacramento, Industrial Blvd. DWR office.

Catch-per-unit effort data, in number per cubic meter of water sampled, for each valid sample are available in Excel with the associated field data by contacting the DWR project lead Jared Frantzich (see contact information at beginning of document).

VIII. Chain of Custody and Sample Handling

Samples are securely packaged to prevent leakage or breakage. All bottles are inspected and verified, and a chain of custody form is filled out with the sample collection time and date, study, site, and number of jars per sample. Signatures are required of both the person responsible for sending the sample package, and the person receiving it. The chain of custody form is signed and sent to the EcoAnalyst contractor with the samples, and the contractor is notified of approximate date of delivery.

Chain of Custody Form

EcoAnalyst Inc. Chain of Custody				Page 1 of 1		
Samples sent from:		West Sacramento		Samples sent to:		Moscow, Idaho
Samples sent by:		DWR, Jared Frantzich		Contract #:		4600009721
Date:				Date:		
Transported By:		UPS		Samples received by:		
Signature:				Signature:		
Requested Analysis:						

	Collection Date	Study	Time	Station	# of Jars	Add Notes
1						
2						
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						
31						
32						
33						
34						
35						
36						
37						
38						
39						

XI. References

A. Taxonomic References

- Abiahy, Bernardo Barroso do, Carlos Eduardo Falavigna da Rocha, and Frank D. Ferrari. 2006. Redescription of *Limnoithona tetraspina* Zhang et Li, 1976 (Copepoda, Cyclopoida) with a discussion of character states shared with the Oithonidae and older cyclopoids. *Invertebrate Zoology* 3(2): 115-135.
- Brooks, John Langdon. 1957. The systematics of North American *Daphnia*. *Memoirs of the Connecticut of Arts & Sciences* Vol. XIII.
- Davis, Charles C. 1955. *The Marine and Fresh-water Plankton*. Michigan State University Press. Michigan.
- Donner, Josef. 1966. Rotifers. Trans. by H. G. S. Wright. Fredrick Warne & Co. Ltd. New York.
- Ferrari, Frank D., and James Orsi. 1984. *Oithona davisae*, new species, and *Limnoithona sinensis* (Burckhardt, 1912) (Copepoda: Oithonidae) from the Sacramento-San Joaquin Estuary, California. *Journal of Crustacean Biology* 4(1): 106-126.
- Light, S.F., Ralph I. Smith, Frank A. Pitelka, Donald P. Abbot, and Frances M. Weesner. 1954. *Intertidal invertebrates of the Central California coast*. University of California Press.
- Mecum, W. Lee. 2007. A Key to the Mysidacea of the Upper San Francisco Estuary. IEP Technical Report 75. 13pp.
- Pennak, Robert W. 1989. *Fresh-water Invertebrates of the United States: Protozoa to Mollusca* 3rd ed. John Wiley and Sons. New York.
- Sars, Georg Ossian. 1918. An account of the Crustacea of Norway : with short descriptions and figures of all the species.: Vol.VI,-Copepoda. Bergen Museum. Bergen.
- Tattersall, W. M. 1932. Contribution to a knowledge of the Mysidacea of California. II. The Mysidacea collected during the survey of San Francisco Bay by the U.S.S. Albatross in 1914. *University of California Publications in Zoology*, 37:301-314.
- Tattersall, W. M. 1951. A review of the Mysidacea of the United States National Museum. *Bulletin of the US National Museum*, 201:1-292.
- Ward, Henry Baldwin, George Chandler Whipple and W. T. Edmondson. 1959. *Fresh-water Biology* 2nd ed. John Wiley and Sons. New York.

B. Program Reports, Publications, and Other Pertinent Literature

- Cranston, P.S., G.M. Benigno, and M.C. Dominguez. 2007. [Hydrobaenus saetheri](#) Cranston, new species, an aestivating, winter-emerging chironomid (Diptera: Chironomidae) from California. Pages 73-79 in *Contributions to the Systematics and Ecology of Aquatic Diptera-A tribute to Ole A. Saether*. T. Andersen, editor. The Caddis Press
- Feyrer, F, T. Sommer, and W. Harrell. 2006. [Managing floodplain inundation for native fish: production dynamics of age-0 splittail in California's Yolo Bypass](#). *Hydrobiology* 573:213-226.
- General Oceanics Inc. *General Oceanics Digital Flowmeter Mechanical and Electronic Operators Manual*. Miami FL. 15 pp.
- Harrell, W.C. and T.R. Sommer. 2003. [Patterns of Adult Fish Use on California's Yolo Bypass Floodplain. Pages 88-93 in P.M. Faber, editor. California riparian systems: Processes and floodplain management, ecology, and restoration](#). 2001 Riparian Habitat and Floodplains Conference Proceedings, Riparian Habitat Joint Venture, Sacramento, California.
- Jassby A.D. and J.E. Cloern 2000. Organic matter sources and rehabilitation of the Sacramento-San Joaquin Delta (California, USA). *Aquatic Conservation: Marine and Freshwater Ecosystems* 10:323-352.
- Kurth, R., and M. Nobriga. 2001 Food Habits of larval splittail. *Interagency Ecological Program Newsletter* 14 (2):40-42
- Lehman, P. W., T. Sommer and L. Rivard. 2008. Phytoplankton primary productivity, respiration, chlorophyll a and species composition in the Yolo Bypass floodplain, California. *Aquatic Ecology* 42:363-378.
- Moyle, P. R. D. Baxter, T. Sommer, T. C. Foin, and S. C. Matern. 2004. Biology and Population Dynamics of Sacramento Splittail (*Pogonichthys macrolepidotus*) in the San Francisco Estuary: a review. *San Francisco Estuary and Watershed Science* [online serial]. Vol. 2, Issue 2 (May 2004), Article 3.

Mueller-Solger, A. B., A. D. Jassby and D. C. Mueller-Navarra. 2002. [Nutritional quality for zooplankton \(Daphnia\) in a tidal freshwater system \(Sacramento-San Joaquin River Delta, USA\)](#). Limnology and Oceanography 47(5):1468-1476.

Schemel, L.E., T.R. Sommer, A.B. Muller-Solger, and W.C. Harrell. 2004. [Hydrologic variability, water chemistry, and phytoplankton biomass in a large floodplain of the Sacramento River, CA, USA](#). Hydrobiologia 513:129-139.

Sommer, T., R. Baxter, and B. Herbold. 1997. [The resilience of splittail in the Sacramento-San Joaquin Estuary](#). Transactions of the American Fisheries Society 126:961-976.

Sommer, T.R., W.C. Harrell, A. Mueller-Solger, B. Tom, and W. Kimmerer. 2004. [Effects of flow variation on channel and floodplain biota and habitats of the Sacramento River, California, USA](#). Aquatic Conservation: Marine and Freshwater Ecosystems 14:247-261.

Sommer, T. R., M. L. Nobriga, W. C. Harrell, W. Batham, and W. J. Kimmerer. 2001. [Floodplain rearing of juvenile Chinook salmon: evidence of enhanced growth and survival](#). Canadian Journal of Fisheries and Aquatic Sciences 58(2):325-333.

Sommer, T, W. Harrell, and M. Nobriga. 2005. [Habitat use and stranding risk of juvenile Chinook salmon on a seasonal floodplain](#). North American Journal of Fisheries Management 25:1493-1504.

Sommer, T., R. Baxter, and F. Feyrer. 2007. [Splittail revisited: how recent population trends and restoration activities led to the "delisting" of this native minnow](#). Pages 25-38 in M.J. Brouder and J.A. Scheuer, editors. Status, distribution, and conservation of freshwater fishes of western North America. American Fisheries Society Symposium 53. Bethesda, Maryland.