

Yolo Bypass Zooplankton Sampling Meta Data – Lynn Takata

Suggested Citation: California Department of Water Resources, Aquatic Ecology Section. Yolo Bypass Zooplankton Sampling Meta Data. 2016.

IEP Aquatic Ecology Section Yolo Bypass Zooplankton Sampling Meta Data Last updated: August 2016

I. Contact Information

Program Manager: Brian Schreier

Lead Contact: Jared Frantzich
Dept. of Water Resources
Division of Environmental Services
3500 Industrial Blvd., West Sacramento, CA.
(209) 942-6088
email: jfrantzi@water.ca.gov

II. Study Element and Objectives

Largely supported by the Interagency Ecological Program (IEP), DWR has operated a fisheries and invertebrate monitoring program in the Yolo Bypass since 1998. The monitoring program has provided a wealth of information regarding the significance of seasonal floodplain habitat to native fishes. Basic objectives of the project are to collect baseline data on lower trophic levels (phytoplankton, zooplankton and insect drift), juvenile and adult fish, hydrology, and physical conditions. As the Yolo Bypass has been identified as a high restoration priority by the US Fish and Wildlife Service and National Marine Fisheries Service biological opinions for Delta Smelt (*Hypomesus transpacificus*) and winter and spring-run Chinook Salmon (*Oncorhynchus tshawytscha*), and by California EcoRestore, these baseline data are critical for evaluating success of future restoration projects. In addition, the data have already served to increase our understanding of the role of the Yolo Bypass in the life history of native fishes, and its ecological function in the San Francisco Estuary. Key findings include: (1) Yolo Bypass is a major factor regulating year class strength of splittail, *Pogonichthys macrolepidotus* (Sommer et al., 1997; Feyrer et al., 2006; Sommer et al., 2007a); (2) Yolo Bypass is a key migration corridor for adult fish of several listed and sport fish (Harrell and Sommer 2003); (3) it is one of the most important regional rearing areas for juvenile Chinook Salmon (Sommer et al., 2001a; 2005); and (4) Yolo Bypass is a source of phytoplankton to the food web of the San Francisco Estuary (Jassby and Cloern 2000; Schemel et al., 2004; Sommer et al., 2004).

The collection of zooplankton is one element of the Aquatic Ecology Section's (AES), Yolo Bypass Fish Monitoring Program's (YBFMP) lower trophic monitoring that is conducted under the IEP umbrella. Zooplankton are an important component in the diet of larval, juvenile, and small adult fishes within the San Francisco Estuary, including Delta Smelt, juvenile Chinook Salmon, Striped Bass, and Sacramento Splittail. The goals of the zooplankton monitoring project are to compare the seasonal variation in species densities and trends within (1) the Sacramento River channel, and (2) the Yolo Bypass, the river's seasonal floodplain.

Key findings to date include: (1) Chinook Salmon sampled in the floodplain contained diets comprised of 90% dipterans and zooplankton, with zooplankton being the dominant prey item in all months (Sommer et al., 2001), (2) laboratory studies showed that the increased chlorophyll-a concentrations in the Yolo Bypass resulted in faster growth rates for the cladoceran *Daphnia magna* as compared to the Sacramento River (Mueller-Solger et al., 2002), and (3) due to high phytoplankton biomass in the spring, the floodplain is suggested to be important in the bottom-up energy transfer through the food web of the San Francisco Estuary (Sommer et al., 2001, Lehman et al., 2007).

III. Study Area and Sample Sites


A. General Information

There are two fixed sampling site locations for this study: (1) Toe Drain of the Yolo Bypass at our rotary screw trap (STTD), and (2) Sacramento River at Sherwood Harbor (SHR). These sites are sampled on an ebb tide on the same day or within one day of one another.

B. Name and Location Information for Zooplankton Sampling Sites

Station	Location	latitude			longitude			Start Year
		degrees	minutes	seconds	degrees	minutes	seconds	
STTD	Yolo Bypass - Screw Trap at Toe Drain	38	21	12.46	121	38	34.71	1999
SHR	Sacramento River at Sherwood Harbor	38	31	56.77	121	31	41.1	1999

Map of Currently Sampled Sites



IV. Period of Record

Zooplankton monitoring began in 1999 and continues through the present. The zooplankton dataset includes the proper sorting, identification, and enumeration of (1) meso-zooplankton (calanoids, cyclopoids, harpacticoids, and cladocerans), (2) microzooplankton and nauplii (rotifers, barnacles, copepod nauplii, cladocera nauplii, and ostracods), and (3) macrozooplankton (mysids, clams, snails, etc.).

V. Sampling Frequency

Following initial pilot years 1999-2001, sampling was conducted at least once monthly during the months of January - June at two sites: One at the rotary screw trap in the Yolo Bypass Toe Drain, and the other at Sherwood Harbor in the Sacramento River. In some years, sampling was conducted weekly during the inundation and draining of the Yolo Bypass floodplain. Since 2011, sampling is conducted at least biweekly (every other week) year-round during non-flooding periods, and weekly during floodplain inundation and drainage events.

Sampling Frequency by Month and Year

Yolo Bypass Screw Trap at Toe Drain (STTD) (150 µm Net)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
1999	0	0*	8*	5	0	0	0	0	0	0	0	0	13
2000	0	4*	3*	0	0	0	0	0	0	0	0	0	7
2001	1	2	2	2	1	1	0	0	0	0	0	0	9
2002	4*	2	3	2	3	2	0	0	0	0	0	0*	16
2003	6*	0	2	2	3*	2	0	0	0	0	0	0	15
2004	2*	2*	2*	2	1	1	0	0	0	0	0	0	10
2005	4	3	4	4	2*	2	0	0	0	0	0	0	19
2006	3*	3*	4*	1*	2*	1	0	0	0	0	0	0	14
2007	0	1	2	1	2	2	0	0	0	0	0	0	8
2008	1	1	2	2	2	2	0	0	0	0	0	0	10
2009	2	2	2	2	2	2	0	0	0	0	0	0	12
2010	4*	3	3	2	2	0	0	0	0	0	0	0*	14
2011	2*	2	2*	3*	2	3	2	2	2	2	3	2	27
2012	2	2	2	2	3	2	2	2	3	3	2	2*	27
2013	4	4	3	3	2	2	2	2	2	3	1	2	30
2014	3	2	2	2	2	2	2	3	2	2	2	3	27
2015	4	4	7	4	3	6	4	4	2	3	2	2	45
Total	42	37	53	40	32	30	12	13	11	13	10	11	303

*Months with overtopping at Fremont Weir.

Sacramento River at Sherwood Harbor (SHR) (150 µm Net)

Year	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
1999	0	0*	8*	5	0	0	0	0	0	0	0	0	13
2000	0	3*	3*	0	0	0	0	0	0	0	0	0	6
2001	1	2	2	2	1	1	0	0	0	0	0	0	9
2002	4*	2	3	2	3	2	0	0	0	0	0	0*	16
2003	6*	1	2	2	3*	2	0	0	0	0	0	0	16
2004	2*	1*	1*	2	1	1	0	0	0	0	0	0	8
2005	4	3	4	4	2*	2	0	0	0	0	0	0*	19
2006	3*	2*	4*	2*	2*	2	0	0	0	0	0	0	15
2007	0	0	2	1	2	1	0	0	0	0	0	0	6
2008	1	2	2	1	2	2	0	0	0	0	0	0*	10
2009	2	2	2	2	2	2	0	0	0	0	0	0	12
2010	4*	4	3	2	2	0	0	0	0	0	0	0*	15
2011	2*	2	2*	3*	2	3	2	2	2	2	3	2	27
2012	2	2	2	2	3	2	2	2	2	3	2	2*	26
2013	5	4	3	3	2	2	2	2	2	3	1	2	31
2014	3	2	2	2	2	2	2	3	2	2	2	3	27
2015	4	4	6	5	4	4	5	5	2	2	2	2	45
Total	43	36	51	40	33	28	13	14	10	12	10	11	301

*Months with overtopping at Fremont Weir.

Number of Sampling Events by Station and by Year

Station	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	Total
STTD	13	7	9	16	15	10	19	15	8	10	12	16	27	26	30	27	45	301
SHR	13	6	9	16	16	9	19	15	6	10	12	16	27	26	31	26	45	302
Total	26	13	18	32	31	19	38	30	14	20	24	32	54	52	61	53	90	604

VI. Field Collection Methods

A simple plankton net is used to capture (1) Calanoids (adult and juvenile copepods), (2) Cyclopoids (adult and juvenile copepods of the genera *Limnithona*, *Oithona*, and *Acanthocyclops*), (3) Cladocerans, (4) Harpacticoids and (5) Microzooplankton and Nauplii (copepod and cladocera nauplii, ostracods, and rotifers).

A. Simple conical plankton net

The plankton net is made of 150 micron mesh net, with a 0.50 m diameter outer mouth (with a General Oceanics Model 2030R flowmeter mounted inside) and 2 meters in length. It tapers to 0.076 m at the cod-end where a polyethylene jar screened with 150 micron mesh collects the organisms. When there is sufficient flow (typically from January – June), Toe Drain samples are collected during the ebb tide from the rotary screw trap anchored in the middle of the channel, and Sacramento River/Sherwood Harbor samples are taken dockside. In the absence of sufficient downstream flow, typically from July-Nov, Sacramento River and Yolo Bypass samples are taken from a boat moving approximately 2-3 mph upstream near the screw trap or dock. Net tow times have varied through the years, with shorter tows occurring with high flows and/or debris loads. Generally, tows have been 5 or 10 minutes long, and tow times are recorded with every sampling event.

All samples are preserved in the field with 10% formalin with Rose Bengal dye to aid in separating organisms from detritus and algae.

Water quality parameters are recorded when the sample is collected. Temperature (C), electrical conductivity (uS/cm), dissolved oxygen (mg/L), and pH are measured using a YSI 556 Multiprobe System. Turbidity is measured from a water sample collected in a glass vial and later analyzed at the office using a Hach 2100Q Portable Turbidimeter. Secchi depth (cm) is also measured. Other factors including tide stage, weather, and trap condition code are also recorded.

VII. Lab Processing Methods

Zooplankton samples are concentrated and retained in the laboratory by pouring them through a sieve screened with 106 micron mesh wire. Excess formalin is rinsed off using tap water, and sample is transferred to 70-80% ETOH before delivery to contractor for taxonomic identification and enumeration: BSA Environmental Services, Inc. (23400 Mercantile Road, Suite 8, Beachwood, Ohio 44122).

A. Sample Analysis Procedure (2015-Current)

1. Each sample is rinsed and filtered through a 153 micron sieve and a 43 micron sieve. The 153 micron sieve removes debris and large non-target organisms. Material on the 43 micron sieve is retained.
2. For mesozooplankton, water is added to the remaining sample to achieve a target of 40-50 organisms per mL, and subsample each 5 times to accumulate a total of 200-250 organisms per sample.
3. For microzooplankton, the sample total volume is readjusted to target 100 organisms per mL, and 3 subsamples are taken to accumulate 300 organisms total per sample.
4. This results in two total volumes and two subsample volumes for each sample; one for mesozooplankton and the other for microzooplankton.
5. The sample is stirred to distribute the organisms homogeneously, and the volume of water added with be recorded as the subsample volume.
6. A sub-sample is extracted with a Hensen Stempel pipette, dispensed into a Ward zooplankton counting wheel, and examined under a compound microscope at a minimum 100x magnification.
7. The sub-sample volume is recorded and zooplankton are enumerated to the lowest taxon possible
8. Mesozooplankton are enumerated by differentiating life stages and species. *Calanoid copepods*, *Acanthocyclops vernalis*, *Oithona* and *Limnoithona* will be identified to species level, and juveniles and adults are recorded separately. *Harpacticoids* are identified to order level, with juveniles and adults combined. *Cladocerans* are identified to the genus level, with juveniles and adults combined. Microzooplankton (i.e. rotifers, barnacles, copepod nauplii, cladocera nauplii, unid. nauplii, ostracods) will be enumerated.
9. If the sample has any suspended sediment, the sediment volume is recorded separately.
10. For selected samples, additional subsamples will be taken to obtain at least 20 adult females of the numerically dominant copepod species unless no copepod is very abundant in the sample. In addition, the analyst also counts egg masses that are clearly identified as coming from the abundant species. Either all of the egg masses, or a subsample of 20 egg masses, are teased apart and the eggs are counted.

B. Calculating Volume of Water Sampled

The number per cubic meter for each zooplankton taxon taken in the net was calculated using the following equation:

$$N = ((C/S)/V)$$

N = the number of a taxon per cubic meter of water sampled

C = the total number of a taxon counted for the sample

S = the total subsample volume

V = the volume of water sampled through the net (m³)

Calculations for volume of water sampled through the net is specific to the General Oceanics Flowmeter model 2030R, and is calculated as follows (General Oceanics Inc.):

$$\frac{(\text{Flowmeter count start} - \text{Flowmeter count end}) \times \text{Rotor Constant}}{999999} \times \frac{\text{Net mouth area}}{4}$$

The rotor constant depends upon which the flowmeter rotor was used during each sampling event, and is identified in the sampling database. Rotor constants are specified in the General Oceanics Flowmeter 2030R manual as:

Standard Speed Rotor Constant = 26,873

Low Speed Rotor Constant R6 = 57,560

Organisms Found in Zooplankton Samples

MICROZOOPLANKTON & NAUPLII	CLADOCERA
Rotifers	Bosmina
Barnacles	Ceriodaphnia
Copepod nauplii	Daphnia
Cladocera nauplii	Chydorus
Unid nauplii	Camptocercus
Ostracods	Scaphloberis
CYCLOPOIDS	Diaphanosoma
Cyclopoid adult	Juvenile Daphnia
Cyclopoid copepodid	Alona
Acanthocyclops vernalis copepodid	Ilyocryptus
Acanthocyclops vernalis adult	Macrothrix
Oithona spp.	HARPACTICIDS
Oithona similis	
Oithona davisae	
Oithona copepodid	
Limnoithona spp.	
Limnoithona tetraspina	
Limnoithona sinensis	
Limnoithona copepodid	
CALANOIDS	
Acartia spp.	
Acartia copepodid	
Diaptomidae	
Diaptomidae copepodid	
Pseudodiaptomus forbesi adult	
Pseudodiaptomus forbesi copepodid	
Eurytemora affinis adult	
Eurytemora affinis copepodid	
Sinocalanus doerrii adult	
Sinocalanus doerrii copepodid	
Acartiella sinensis	
Acartiella copepodid	
Tortanus spp.	
Tortanus copepodid	
Osphranticum labronectum	
Osphranticum copepodid	

VIII. Data Management and Quality Assurance/Quality Control

A. Field Data

Field data are collected and recorded onto datasheets by DWR personnel. These data are then entered monthly by DWR personnel into an Access database. Field data are reviewed monthly for accuracy and completeness. Annually, after all samples are processed for the year, lab data are reviewed for accuracy and completeness.

B. Field Datasheet

Paper datasheets are digitized and archived in binders that are stored at the West Sacramento, Industrial Blvd. DWR office.

Field Datasheet

LOWER TROPHIC SAMPLING – YOLO BYPASS STUDY									
2015/2016									
Location: _____		Crew: _____		Date: _____		Time: _____		pH: _____	
Secchi Depth: _____ m		Water Temp: _____ °C		Weather: _____		Turb: _____ NTU		DO: _____	
Vial #: _____		SpCnd: _____		Cnd (EC): _____					
Light Attenuation:									
Surface Irradiance (in air avg):		Subsurface Irradiance (in water avg) (~75%, ~50%, ~25%, ~1%):							
LL-COR Calibration -143.27 (in air) -232.10 (in water)		0.75 = _____ μmol		0.50 = _____ μmol		0.25 = _____ μmol		0.01 = _____ μmol	
Depth: _____ m		① _____ μmol		③ _____ μmol		Depth: _____ m		④ _____ μmol	
② _____ μmol		Depth: _____ m		④ _____ μmol					
Drift Sample :									
Start Time: _____		Stop Time: _____		Set Time: _____ min		Condition Code: _____			
Flow Meter: _____		Flow : _____		Start Meter: _____		End Meter: _____			
Regular or Low Speed		*For low speed, record initial meter reading in "end meter" box							
Comments: _____									
Egg & Larval Fish Sample :									
1st Start Time: _____		1st Stop Time: _____		2nd Start Time: _____		2nd Stop Time: _____		Set Time: _____ min	
Flow Meter: _____		Flow : (Mid-West) Start Meter: _____		End Meter: _____		(Near-West) Start Meter: _____		End Meter: _____	
Regular or Low Speed		*For low speed, record initial meter reading in "end meter" box		(Mid-East) Start Meter: _____		End Meter: _____		(Near-East) Start Meter: _____	
Comments: _____		(Near-East) Start Meter: _____		End Meter: _____					
Zooplankton Sample :									
Start Time: _____		150 Stop Time: _____		50 Stop Time: _____		150 Set Time: _____ min		50 Set Time: _____ min	
Flow Meter: _____		Flow : _____		150μm: Start Meter: _____		End Meter: _____		50μm: Start Meter: _____	
Regular or Low Speed		*For low speed, record initial meter reading in "end meter" box		End Meter: _____					
Comments: _____		50μm: Start Meter: _____		End Meter: _____					
Chlorophyll Sample :									
Yes _____		No _____		Replicate _____		Time: _____		Filtered: 500mL OR 250mL	
Phytoplankton Sample :									
Yes _____		No _____		60 mL Amber Bottle w/ Lugol Solution					
Entered by : _____ Date: _____ Checked by : _____ Date: _____									

C. Taxonomic Data

Taxonomic results are received via email from the contractor and entered into the AES Access database by DWR personnel. Electronic copies of results for taxonomic analyses are archived on DWR/AES Network drives. Hard copies are printed and stored in binders at the West Sacramento, Industrial Blvd. DWR office.

Catch-per-unit effort data, in number per cubic meter of water sampled, for each valid sample are available in Excel with the associated field data by contacting the DWR project lead Jared Frantzich (see contact information at beginning of document).

IX. Chain of Custody and Sample Handling

Samples are securely packaged to prevent leakage or breakage. All bottles are inspected and verified, and a chain of custody form is filled out with the sample collection time and date, study, site, and number of jars per sample. Signatures are required of both the person responsible for sending the sample package, and the person receiving it. The chain of custody form is signed and sent to the BSA contractor with the samples, and the contractor is notified of approximate date of delivery.

Example Chain of Custody Form

Page 1 of 2

BSA Environmental Services, Inc.
Chain of Custody

Samples sent from: West Sacramento	Samples sent to: Beachwood, Ohio
Samples sent by: DWR, Jared Frantzich	Contract #: 4600011015
Date: 8/16/2016	Date:
Transported By: UPS	Samples received by: Dr. John R. Beaver
Signature: <i>Jared Frantzich</i> 8/15/16	Signature:
Requested Analysis: Zooplankton	

	Station	Date	ZoopNetType	Time	# of Jars	Comments
✓ 1	SHR	3/30/2016	50	8:28	1	
✓ 2	SHR	3/30/2016	150	8:28	1	
✓ 3	SHR	4/7/2016	50	10:05	1	
✓ 4	SHR	4/7/2016	150	10:05	1	
✓ 5	SHR	4/18/2016	50	10:55	1	
✓ 6	SHR	4/18/2016	150	10:55	1	
✓ 7	SHR	5/5/2016	50	8:38	1	
✓ 8	SHR	5/5/2016	150	8:38	1	
✓ 9	SHR	5/20/2016	50	9:37	1	
✓ 10	SHR	5/20/2016	150	9:37	1	
✓ 11	SHR	5/31/2016	50	9:15	1	
✓ 12	SHR	5/31/2016	150	9:15	1	
✓ 13	SHR	6/13/2015	50	9:17	1	
✓ 14	SHR	6/13/2015	150	9:17	1	
✓ 15	SHR	6/29/2016	50	9:00	1	
✓ 16	SHR	6/29/2016	150	9:00	1	
✓ 17	SHR	7/6/2016	150	8:57	1	
✓ 18	STTD	3/30/2016	50	10:51	1	
✓ 19	STTD	3/30/2016	150	10:51	2 ✓	
✓ 20	STTD	4/8/2016	50	9:58	1	
✓ 21	STTD	4/8/2016	150	9:58	1	
✓ 22	STTD	4/19/2016	50	9:19	1	
✓ 23	STTD	4/19/2016	150	9:19	1	
✓ 24	STTD	5/4/2016	50	10:59	1	
✓ 25	STTD	5/4/2016	150	10:59	1	
✓ 26	STTD	5/19/2016	50	10:08	1	
✓ 27	STTD	5/19/2016	150	10:08	1	
✓ 28	STTD	6/1/2016	50	10:04	1	
✓ 29	STTD	6/1/2016	150	10:04	1	
✓ 30	STTD	6/14/2016	50	9:40	1	
✓ 31	STTD	6/14/2016	150	9:40	3 ✓	
✓ 32	STTD	6/30/2016	50	10:40	1	
✓ 33	STTD	6/30/2016	150	10:40	1	
✓ 34	BL5	6/30/2016	150	11:30	2	
✓ 35	LIB	6/30/2016	150	12:15	1	

Example Chain of Custody Form (Continued)

Samples sent to: Beachwood, Ohio
Contract #: 4600011015
Date:
Samples received by: Dr. John R. Beaver
Signature: *[Handwritten Signature]* 8/18/16 3:30

[illegible]

XI. References

A. Taxonomic References

- Abiahy, Bernardo Barroso do, Carlos Eduardo Falavigna da Rocha, and Frank D. Ferrari. 2006. Redescription of *Limnoithona tetraspina* Zhang et Li, 1976 (Copepoda, Cyclopoida) with a discussion of character states shared with the Oithonidae and older cyclopoids. *Invertebrate Zoology* 3(2): 115-135.
- Brooks, John Langdon. 1957. The systematics of North American *Daphnia*. *Memoirs of the Connecticut of Arts & Sciences* Vol. XIII.
- Davis, Charles C. 1955. *The Marine and Fresh-water Plankton*. Michigan State University Press. Michigan.
- Donner, Josef. 1966. Rotifers. Trans. by H. G. S. Wright. Fredrick Warne & Co. Ltd. New York.
- Ferrari, Frank D., and James Orsi. 1984. *Oithona davisae*, new species, and *Limnoithona sinensis* (Burckhardt, 1912) (Copepoda: Oithonidae) from the Sacramento-San Joaquin Estuary, California. *Journal of Crustacean Biology* 4(1): 106-126.V
- Light, S.F., Ralph I. Smith, Frank A. Pitelka, Donald P. Abbot, and Frances M. Weesner. 1954. *Intertidal invertebrates of the Central California coast*. University of California Press.
- Mecum, W. Lee. 2007. A Key to the Mysidacea of the Upper San Francisco Estuary. IEP Technical Report 75. 13pp.
- Pennak, Robert W. 1989. *Fresh-water Invertebrates of the United States: Protozoa to Mollusca* 3rd ed. John Wiley and Sons. New York.
- Sars, Georg Ossian. 1918. An account of the Crustacea of Norway : with short descriptions and figures of all the species.: Vol.VI,-Copepoda. Bergen Museum. Bergen.
- Tattersall, W. M. 1932. Contribution to a knowledge of the Mysidacea of California. II. The Mysidacea collected during the survey of San Francisco Bay by the U.S.S. Albatross in 1914. *University of California Publications in Zoology*, 37:301-314.
- Tattersall, W. M. 1951. A review of the Mysidacea of the United States National Museum. *Bulletin of the US National Museum*, 201:1-292.
- Ward, Henry Baldwin, George Chandler Whipple and W. T. Edmondson. 1959. *Fresh-water Biology* 2nd ed. John Wiley and Sons. New York.

B. Program Reports, Publications, and Other Pertinent Literature

- Cranston, P.S., G.M. Benigno, and M.C. Domingeuz. 2007. [*Hydrobaenus saetheri* Cranston, new species, an aestivating, winter-emerging chironomid \(Diptera: Chironomidae\) from California](#). Pages 73-79 in Contributions to the Systematics and Ecology of Aquatic Diptera-A tribute to Ole A. Saether. T. Andersen, editor. The Caddis Press
- Feyrer, F, T. Sommer, and W. Harrell. 2006. [Managing floodplain inundation for native fish: production dynamics of age-0 splittail in California's Yolo Bypass](#). *Hydrobiology* 573:213-226.
- General Oceanics Inc. General Oceanics Digital Flowmeter Mechanical and Electronic Operators Manual. Miami FL. 15 pp.
- Harrell, W.C. and T.R. Sommer. 2003. [Patterns of Adult Fish Use on California's Yolo Bypass Floodplain. Pages 88-93 in P.M. Faber, editor. California riparian systems: Processes and floodplain management, ecology, and restoration](#). 2001 Riparian Habitat and Floodplains Conference Proceedings, Riparian Habitat Joint Venture, Sacramento, California.
- Jassby A.D. and J.E. Cloern 2000. Organic matter sources and rehabilitation of the Sacramento-San Joaquin Delta (California, USA). *Aquatic Conservation: Marine and Freshwater Ecosystems* 10:323-352.
- Kurth, R., and M. Nobriga. 2001 Food Habits of larval splittail. *Interagency Ecological Program Newsletter* 14 (2):40-42

- Lehman, P. W., T. Sommer and L. Rivard. 2008. Phytoplankton primary productivity, respiration, chlorophyll a and species composition in the Yolo Bypass floodplain, California. *Aquatic Ecology* 42:363-378.
- Moyle, P. R. D. Baxter, T. Sommer, T. C. Foin, and S. C. Matern. 2004. Biology and Population Dynamics of Sacramento Splittail (*Pogonichthys macrolepidotus*) in the San Francisco Estuary: a review. *San Francisco Estuary and Watershed Science* [online serial]. Vol. 2, Issue 2 (May 2004), Article 3.
- Mueller-Solger, A. B., A. D. Jassby and D. C. Mueller-Navarra. 2002. [Nutritional quality for zooplankton \(Daphnia\) in a tidal freshwater system \(Sacramento-San Joaquin River Delta, USA\)](#). *Limnology and Oceanography* 47(5):1468-1476.
- Schemel, L.E., T.R. Sommer, A.B. Muller-Solger, and W.C. Harrell. 2004. [Hydrologic variability, water chemistry, and phytoplankton biomass in a large floodplain of the Sacramento River, CA, USA](#). *Hydrobiologia* 513:129-139.
- Sommer, T., R. Baxter, and B. Herbold. 1997. [The resilience of splittail in the Sacramento-San Joaquin Estuary](#). *Transactions of the American Fisheries Society* 126:961-976.
- Sommer, T.R., W.C. Harrell, A. Mueller-Solger, B. Tom, and W. Kimmerer. 2004. [Effects of flow variation on channel and floodplain biota and habitats of the Sacramento River, California, USA](#). *Aquatic Conservation: Marine and Freshwater Ecosystems* 14:247-261.
- Sommer, T. R., M. L. Nobriga, W. C. Harrell, W. Batham, and W. J. Kimmerer. 2001. [Floodplain rearing of juvenile Chinook salmon: evidence of enhanced growth and survival](#). *Canadian Journal of Fisheries and Aquatic Sciences* 58(2):325-333.
- Sommer, T, W. Harrell, and M. Nobriga. 2005. [Habitat use and stranding risk of juvenile Chinook salmon on a seasonal floodplain](#). *North American Journal of Fisheries Management* 25:1493-1504.
- Sommer, T., R. Baxter, and F. Feyrer. 2007. [Splittail revisited: how recent population trends and restoration activities led to the "delisting" of this native minnow](#). Pages 25-38 in M.J. Brouder and J.A. Scheuer, editors. *Status, distribution, and conservation of freshwater fishes of western North America*. American Fisheries Society Symposium 53. Bethesda, Maryland.