



Peer Reviewed

Title:

The Role of Tidal Marsh Restoration in Fish Management in the San Francisco Estuary

Journal Issue:

[San Francisco Estuary and Watershed Science, 12\(1\)](#)

Author:

[Herbold, Bruce](#), U.S. Environmental Protection Agency, retired
[Baltz, Donald M.](#), Dept. of Oceanography and Coastal Sciences, Louisiana State University
[Brown, Larry](#), U.S. Geological Survey
[Grossinger, Robin](#), San Francisco Estuary Institute
[Kimmerer, Wim](#), Romberg Tiburon Center, San Francisco State University
[Lehman, Peggy](#), California Department of Water Resources
[Simenstad, Charles Si](#), University of Washington
[Wilcox, Carl](#), California Dept. Fish and Wildlife
[Nobriga, Matthew](#), U.S. Fish and Wildlife Service

Publication Date:

2014

Publication Info:

San Francisco Estuary and Watershed Science

Permalink:

<http://www.escholarship.org/uc/item/1147j4nz>

Acknowledgements:

The authors thank the Delta Science Program, the University of California, Davis' Center for Aquatic Biology and Aquaculture, and the California–Nevada Chapter of the American Fisheries Society for their funding and logistical support of this symposium.

Author Bio:

Wetland Ecosystem Team

Bay-Delta Region

Keywords:

Tidal Marsh, Restoration, Productivity, Dispersion, Ecology

Local Identifier:

jmie_sfews_18827



Abstract:

Tidal marsh restoration is an important management issue in the San Francisco Estuary (estuary). Restoration of large areas of tidal marsh is ongoing or planned in the lower estuary (up to 6,000 ha, Callaway et al. 2011). Large areas are proposed for restoration in the upper estuary under the Endangered Species Act biological opinions (3,237 ha) and the Bay Delta Conservation Plan (26,305 ha). In the lower estuary, tidal marsh has proven its value to a wide array of species that live within it (Palaima 2012). In the Sacramento–San Joaquin Delta (Delta), one important function ascribed to restoration of freshwater tidal marshes is that they make large contributions to the food web of fish in open waters (BDCP 2013). The Ecosystem Restoration Program ascribed a suite of ecological functions to tidal marsh restoration, including habitat and food web benefits to native fish (CDFW 2010). This background was the basis for a symposium, *Tidal Marshes and Native Fishes in the Delta: Will Restoration Make a Difference?* held at the University of California, Davis, on June 10, 2013. This paper summarizes conclusions the authors drew from the symposium.

Copyright Information:



Copyright 2014 by the article author(s). This work is made available under the terms of the Creative Commons Attribution 3.0 license, <http://creativecommons.org/licenses/by/3.0/>



eScholarship
University of California

eScholarship provides open access, scholarly publishing services to the University of California and delivers a dynamic research platform to scholars worldwide.

The Role of Tidal Marsh Restoration in Fish Management in the San Francisco Estuary

Bruce Herbold¹, Donald M. Baltz², Larry Brown³, Robin Grossinger⁴, Wim Kimmerer⁵, Peggy Lehman⁶, Peter B. Moyle⁷, Matt Nobriga⁸, and Charles A. Simenstad⁹

INTRODUCTION

Tidal marsh restoration* is an important management issue in the San Francisco Estuary (estuary). Restoration of large areas of tidal marsh is ongoing or planned in the lower estuary (up to 6,000 ha, Callaway et al. 2011). Large areas are proposed for restoration in the upper estuary under the Endangered Species Act biological opinions (3,237 ha) and the Bay Delta Conservation Plan (26,305 ha). In the lower estuary, tidal marsh has proven its value to a wide array of species that live within it (Palaima 2012). In the Sacramento–San Joaquin Delta (Delta), one important function ascribed to restoration of freshwater tidal marshes is that they make large contributions to the food web of fish in open waters

(BDCP 2013). The Ecosystem Restoration Program ascribed a suite of ecological functions to tidal marsh restoration, including habitat and food web benefits to native fish (CDFW 2010). This background was the basis for a symposium, *Tidal Marshes and Native Fishes in the Delta: Will Restoration Make a Difference?* held at the University of California, Davis, on June 10, 2013. This paper summarizes conclusions the authors drew from the symposium.

CONSENSUS CONCLUSIONS

From the scientific work done in the estuary and elsewhere we conclude:

1. Restoration of tidal marshes benefits many fish, mammals, and birds. These benefits can be extremely important for growth and survival of individuals of desirable species on site. Site location of restored marshes will determine which species will use them. Site-specific design is, therefore, required to support targeted species and to reduce the effects of invasive species. Important design considerations include area, elevations, residence time, extent of edge and channels, the nature of adjacent habitats, and connectivity with adjacent habitats.

* Restoration as used here implies a reversal of impaired ecological features and processes in order to support desired species of wildlife, not a return to historic conditions.

1 Corresponding author: U.S. Environmental Protection Agency, retired; bherbold@gmail.com.

2 Louisiana State University, Baton Rouge, LA USA

3 U.S. Geological Survey, Menlo Park, CA USA

4 San Francisco Estuary Institute, Oakland, CA USA

5 San Francisco State University, San Francisco, CA USA

6 California Department of Water Resources, Sacramento, CA USA

7 University of California Davis, Davis CA USA

8 U.S. Fish and Wildlife Service, CA, USA

9 University of Washington, WA USA

2. Movement of plankton from a tidal marsh (beyond the immediate area of tidal exchange) is likely to be limited and to decrease strongly with distance. Even under ideal circumstances, plankton in water discharged from tidal marsh cannot greatly affect the standing crop of plankton in large, deep channels. Feeding by clams and other introduced species can further reduce contributions of marsh plankton to open-water foodwebs.
3. Large areas with diverse physical structure will enhance habitat diversity and help meet the various needs of targeted species. No quantitative guidelines exist to relate restoration extent to functional contributions at the population scale, but a good starting point is to focus on areas large enough to support tidal channels of diverse size and density similar to natural Delta tidal marshes. Diverse habitat types provide benefits to an array of desirable species at multiple life stages.
4. Effective tidal marsh planning requires a landscape-level perspective at a decadal, or greater, scale. Large-scale construction of tidal marsh will change tidal dynamics and alter the tidal inundation regime over a broad area. Sea level rise and inundation of Delta islands will also change tidal dynamics, as will changes in timing or quantity of freshwater flow that results from management or climate change. Tidal wetland design must plan for future tidal and flow regimes.
5. Information gaps about functions and processes in Delta tidal marshes are large but can be filled by designing restoration projects as experiments. In particular, larger restoration areas may produce changes in system response that are large enough to be detected. Planning for new tidal marsh should use site-specific modeling to develop realistic expectations and testable hypotheses, incorporate experimental design to test hypotheses, actively investigate ecological mechanisms that develop in new environments, and contribute toward landscape-level ecological models.

Tidal wetlands elsewhere make broad, multi-faceted contributions to fish habitat, productivity and resilience. However, the present Delta has comparatively little tidal marsh (less than 5% of the historical extent) and so its role is little understood. Experience from previous restoration efforts throughout the estuary, both intentional and accidental, can guide future work. The consensus of the group was that restoration of tidal marsh should proceed both boldly and carefully. Restoration should be accompanied by studies that fill crucial information gaps to help navigate the environmental changes expected in the coming decades.

SELECTED FINDING

Tidal marsh was the dominant component of the primeval Delta (over 90% of its area) and was probably key to historical fish productivity, now largely lost. Other elements of the landscape—including the natural hydrograph, floodplains, sediment supply, and slough networks—are also greatly altered. These alterations and abundant alien species preclude a return to the original Delta. Climate change, earthquakes, and future species invasions will further alter the Delta. Creation and management of tidal marshes can help protect species and ecosystem services that humans value.

Historical records and maps reveal an intricate mosaic of diverse habitats dispersed across three main Delta regions: a floodplain region off the Sacramento River, a meandering channel region from the San Joaquin River and a tidal region where the rivers join before flowing into Suisun Bay (Whipple et al. 2012). Lakes and marshes, riparian forests and seasonal wetlands, and other landscape forms were inundated to different depths and durations during different seasons and years, providing a diverse portfolio of aquatic habitats. Overall, wetland area exceeded open-water area by about 14:1; today, wetland area is less than open water area by a ratio of 1:6, an 80-fold switch in dominant habitat types (Whipple et al. 2012).

Shallow areas like those of the ancient San Francisco Estuary are nurseries for fish in estuaries along the Gulf Coast, the Pacific Northwest, and Chesapeake Bay. Small fish use edges of wetlands to feed and to avoid predation by larger fish (Baltz et al. 1993, 1998). Fish-eating wading birds enhance nursery function by preying on larger fish, thus reducing the risk of predation for small fish. The nursery value of a wetland for a particular species is affected by both accessibility and areal extent. In Louisiana, marsh value is affected by both edge and area. In early stages of degradation, shrinking wetlands retain their value for young fish because the amount of edge increases as wetlands are initially fragmented, which increases fish access (Chesney et al. 2000). On the other hand, harvest-per-hectare of commercial shrimp increases with marsh area, presumably because shrimp are not restricted to the edge (Turner 1977). Thus, the processes that benefit wetland species differ strongly from species to species. Black rails and clapper rails in the lower estuary have a minimum marsh size of about 50 ha and clapper rails have an optimum patch shape with minimum edge-to-area ratio (Spautz and Nur 2002; Liu et al. 2012). Thus, it is crucial to understand marsh characteristics important to each species when size, location, and configuration of new tidal marshes are determined.

Reclaiming tidal wetlands from salt harvest, military use, and agriculture has been a major effort in the estuary for the last 40 years and has improved our understanding of tidal marsh processes. A 2003 summary of the value of tidal wetlands to native fishes found large gaps in knowledge and many unfounded assumptions about tidal marsh function for fishes (Brown 2003). Much knowledge has been gained since 2003 and a revised summary of the current knowledge and knowledge gaps is expected in 2014 (L. Brown, USGS, pers. comm., 2014). Most knowledge has been garnered incidental to restoration activity, rather than as an integrated part of it. For example, isotope studies have been conducted in several tidal marshes with different restoration histories along the Napa River. These studies showed that fish draw much of their nutrition from upstream sources

during wet periods, but that nutrition comes largely from tidal marsh and marine sources when river flow declines and tidal influence increases (Howe and Simenstad 2007, 2011). Three broad themes have emerged about fish use of restored tidal marsh:

1. Food web pathways for fish within a marsh are largely detritus-based, rather than phytoplankton-based (Howe and Simenstad 2007, 2011).
2. The vegetated edge is important for small fish foraging and predator avoidance (Gewant and Bollens 2012).
3. Newly-constructed marshes are rapidly occupied by fish and their prey; new marshes provide habitat and food web support comparable to reference sites (Howe and Simenstad 2007, 2011; Cohen and Bollens 2008).

In the modern San Francisco Estuary, tidal wetlands can be important habitats for many fishes, but likely will have little effect on the export of food available to fish at any significant distance. Measured flux of organic material into and out of Liberty Island (flooded in 1997, now tidal marsh and open water) suggests that little of the productivity that supports pelagic food webs on-site is exported (Lehman et al. 2010). For small fishes like delta smelt, the value of on-site productivity is presumably enhanced by low populations of invasive clams, aquatic plants, and predators in Liberty Island (Sommer and Mejia 2013) and similar areas in Suisun Marsh (P. Moyle, UCD, unpublished data). Seasonal floods bring riverine materials into Liberty Island, but daily tidal action seems not to move much material off-site; data are lacking on export of material that may occur during occasional large-scale flood events. Tidal wetland channels can facilitate phytoplankton growth and accumulation if they are shallow and clear enough that light penetrates most of the water column. Long residence time allows buildup of high biomass, which can fuel further phytoplankton and zooplankton development. Benthic algae can be important parts of primary productivity in shallow or low-turbidity areas. Conversely, the grazing effects of clams are

heightened in shallow water with long residence time (Lucas and Thompson 2012). Therefore, optimizing tidal wetland benefits to fish requires a balance between water depth and residence time to promote planktonic and benthic algal growth while minimizing clam effects. Such balancing requires site-specific design considerations and improved understanding of factors that affect clam abundance.

Restored tidal wetlands are unlikely to have much effect on food webs in the upper estuary's open waters. The shallow depth and small volume of water on tidal wetlands compared to the vast volume of open water in Delta channels and Suisun Bay means that flux of wetland phytoplankton and zooplankton would be inconsequential to pelagic food webs. We are unaware of reports from the worldwide literature in which substantial quantities of zooplankton are exported from marshes to open waters, whereas several studies show net import of zooplankton to fish consumption on site.

Tidal wetland restoration without analysis of processes in the developing ecosystem and in the landscape overall, wastes opportunities to learn from ongoing projects and to improve design of future projects. For example, breaching of dikes at Blacklock in Suisun Marsh was accompanied by little effort to study evolution of the site, so insights to guide future restoration are limited. If levee work does not keep pace with sea level rise, more of Suisun Marsh may become tidal than the amount considered in the Suisun Marsh Plan (USBR 2011). Inundation of large parts of Suisun Marsh would reduce tidal energy entering the Delta and change inundation patterns (and salinity) at other tidal wetland sites. Similarly, inundation of lands in the Delta will alter tidal dynamics throughout the Delta. Thus, studies are needed on restored sites, in areas adjacent to restored sites, and in areas that are affected by changes in hydrodynamics resulting from the restored sites. In short, landscape-level analyses of restoration effects are essential.

Tidal marsh restoration outcomes are site-specific, in that different sites will support different species

and functions based on location, elevation, adjacent habitats, and degree of hydrodynamic connectivity. Tidal marsh restoration can benefit a wide variety of birds, mammals, and plants but to support target fish populations, tidal wetland restoration must target sites that can be accessed by desired fish species and that are minimally affected by invasive species. In the western Delta, the reach from Suisun Marsh to Liberty Island may provide an opportunity for landscape-scale restoration and increase the habitat suitability for a variety of native fish (Moyle et al. 2012; Hanak et al. 2013). Integrated, multi-purpose designs have been developed for some specific sites, including McCormack-Williamson Tract, Dutch Slough, and Prospect Island.

Achieving successful restoration outcomes is severely constrained by many external factors including: alien species, Delta water management, sea level rise, climate change, sediment supply, and contaminants. Alien species and altered habitats dominate most of the Delta and have profound effects on the aquatic ecosystem. The value of tidal wetland restoration to native species will be greatest where aliens are less abundant or where conditions can be altered to reduce their effects. Climate change, sea level rise, and invasive species will require knowledge and flexibility if desirable traits are to be restored to estuarine ecosystems. Early restoration efforts must be approached as experiments in management that will guide later efforts, and be integrated over the entire estuary. We must increase our knowledge of the trajectories of restoration if we are to achieve our goals and adequately respond to future challenges.

ACKNOWLEDGEMENTS

The authors thank the Delta Science Program, the University of California, Davis' Center for Aquatic Biology and Aquaculture, and the California–Nevada Chapter of the American Fisheries Society for their funding and logistical support of this symposium.

REFERENCES

- Baltz DM, Rakocinski C, Fleeger JW. 1993. Microhabitat use by marsh-edge fishes in a Louisiana estuary. *Env Biol Fish* 36:109–126.
- Baltz DM, Fleeger JW, Rakocinski C, McCall JN. 1998. Food, density, and microhabitat: factors affecting growth and recruitment potential of juvenile saltmarsh fishes. *Env Biol Fish* 53:89–103.
- Brown LR. 2003. Will tidal wetland restoration enhance populations of native fishes? *San Fran Est Water Sci* [Internet]. [cited 03 March 2014]; (1)1. Available from: <http://escholarship.org/uc/item/2cp4d8wk>.
- Callaway JC, Parker VT, Vasey MC, Schile LM, Herbert ER. 2011. Tidal wetland restoration in San Francisco Bay: history and current issues. *San Fran Est Water Sci* [Internet]. [cited 03 March 2014]; 9(3). Available from: <http://www.escholarship.org/uc/item/5dd3n9x3>.
- Chesney EJ, Baltz DM, Thomas RG. 2000. Issues and trends in estuarine and coastal fisheries and fish habitat in Louisiana: perspectives from a fish's eye view. *Ecol App* 10(2):350–366.
- Cohen SE, Bollens SM. 2008. Diet and growth of non-native Mississippi silversides and yellowfin gobies in restored and natural wetlands in the San Francisco Estuary. *Mar Ecol Prog Ser* 368:241–254.
- [CDFW] California Department of Fish and Wildlife. 2010. Ecosystem Restoration Program conservation strategy for Stage 2 implementation, Sacramento–San Joaquin Delta ecological management zone. Sacramento (CA): The Resources Agency.
- Gewant D, Bollens SM. 2012. Fish assemblages of interior tidal marsh channels in relation to environmental variables in the upper San Francisco Estuary. *Env Biol Fish* 94(2):483–499.
- Hanak E, Lund J, Durand J, Fleenor W, Gray B, Medellín-Azuara J, Mount J, Moyle P, Phillips C, Thompson B. 2013. Stress relief prescriptions for a healthier Delta ecosystem. San Francisco (CA): Public Policy Institute of California. Available from: <http://www.ppic.org/main/publication.asp?i=1051>.
- Howe ER, Simenstad CA. 2007. Characterizing restoration trajectories through food web linkages in San Francisco Bay's estuarine marshes: a manipulative translocation experiment. *Mar Ecol Prog Ser* 351:65–76.
- Howe ER, Simenstad CA. 2011. Isotopic determination of food web origins in restoring and ancient estuarine wetlands of the San Francisco Bay and Delta. *Estuar Coasts* 34:597–617.
- Lehman PW, Mayr S, Mecum L, Enright C. 2010. The freshwater marsh at Liberty Island, CA, was both a source and sink of inorganic and organic matter to the San Francisco Estuary. *Aquat Ecol* 44:359–372.
- Liu L, Wood J, Nur N, Salas L, Jongsomjit D. 2012. California clapper rail (*Rallus longirostris obsoletus*) population monitoring: 2005–2011. Technical report to the California Department of Fish and Game. San Rafael (CA): PRBO Conservation Science.
- Lucas LV, Thompson JK. 2012. Changing restoration rules: exotic bivalves interact with residence time and depth to control phytoplankton productivity. *Ecosphere* 3(12):117. Available from: <http://dx.doi.org/10.1890/ES12-00251.1>.
- Moyle PB, Bennett W, Durand J, Fleenor W, Gray B, Hanak E, Lund J, Mount J. 2012. Where the wild things aren't: making the Delta a better place for native species. San Francisco (CA): Public Policy Institute of California. Available from: <http://www.ppic.org/main/publication.asp?i=1025>.

SAN FRANCISCO ESTUARY & WATERSHED SCIENCE

Palaima A, editor. 2012. Ecology, conservation and restoration of tidal marshes San Francisco Estuary. Berkeley (CA): UC Press. 265 p.

Sommer T, Mejia F. 2013. A place to call home: a synthesis of delta smelt habitat in the upper San Francisco Estuary. San Fran Est Water Sci [Internet]. [cited 03 March 2014]; 11(2). Available from: <http://www.escholarship.org/uc/item/32c8t244>.

Spautz H, Nur N. 2002. Distribution and abundance in relation to habitat and landscape features and nest site characteristics of California black rail (*Laterallus jamaicensis coturniculus*) in the San Francisco Bay Estuary. A report to the U.S. Fish and Wildlife Service, Coastal Program. Petaluma (CA): PRBO Conservation Science. Available from: http://www.prbo.org/cms/docs/wetlands/BLRA_PRBO_Mar2002.pdf.

Turner RE. 1977. Intertidal vegetation and commercial yields of Penaeid shrimp. Trans Am Fish Soc. 106(5):411–416.

[USBR] U.S. Bureau of Reclamation 2011. Suisun Marsh habitat management, preservation and restoration plan. Available from: http://www.usbr.gov/mp/nepa/nepa_projdetails.cfm?Project_ID=781.

Whipple AA, Grossinger RM, Rankin D, Stanford B, Askevold RA. 2012. Sacramento–San Joaquin Delta historical ecology investigation: exploring pattern and process. Prepared for the California Department of Fish and Wildlife’s Ecosystem Restoration Program. A Report of SFEI–ASC’s Historical Ecology Program, Publication #672. Richmond (CA): San Francisco Estuary Institute–Aquatic Science Center. Available from: <http://www.sfei.org/DeltaHEStudy>.